




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/13, C07K 15/06 C12P 21/08, C12N 5/10 A61K 39/395, C12N 15/62	A1	(11) International Publication Number: WO 94/04678 (43) International Publication Date: 3 March 1994 (03.03.94)
(21) International Application Number: PCT/EP93/02214 (22) International Filing Date: 18 August 1993 (18.08.93) (30) Priority data: 92402326.0 21 August 1992 (21.08.92) EP (34) Countries for which the regional or international application was filed: 93401310.3 21 May 1993 (21.05.93) EP (34) Countries for which the regional or international application was filed: AT et al. (71)(72) Applicants and Inventors: CASTERMAN, Cécile [BE/BE]; HAMERS, Raymond [BE/BE]; Vijversweg 15, B-1640 Sint-Genesius-Rode (BE).		(74) Agents: GUTMANN, Ernest et al.; Ernest Gutmann - Yves Plasseraud S.A., 67, boulevard Haussmann, F-75008 Paris (FR). (81) Designated States: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KP, KR, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: IMMUNOGLOBULINS DEVOID OF LIGHT CHAINS



A schematic representation of Camel immunoglobulins IgG_1 , the putative IgG_2 and IgG_3 . The large (Pro-X)₁₂ of the putative IgG_2 molecule can be modelled into a 6 aa repeat

(57) Abstract

The invention relates to isolated immunoglobulin, characterized in that it comprises two heavy polypeptide chains sufficient for the formation of a complete antigen binding site or several antigen binding sites, this immunoglobulin being further devoid of light polypeptide chains.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

IMMUNOGLOBULINS DEVOID OF LIGHT CHAINS

The invention relates to new isolated immunoglobulins which are devoid of light polypeptide chains. These immunoglobulins do not consist in the degradation products of immunoglobulins composed of both heavy polypeptide and light polypeptide chains but to the contrary, the invention defines a new member of the family of the immunoglobulins, especially a new type of molecules capable of being involved in the immune recognition. Such immunoglobulins can be used for several purposes, especially for diagnosis or therapeutical purposes including protection against pathological agents or regulation of the expression or activity of proteins.

Up to now the structure proposed for immunoglobulins consists of a four-chain model referring to the presence of two identical light polypeptide chains (light chains) and two identical heavy polypeptide chains (heavy chains) linked together by disulfide bonds to form a y- or T-shaped macromolecules. These chains are composed of a constant region and a variable region, the constant region being subdivided in several domains. The two heavy polypeptide chains are usually linked by disulphide bounds in a so-called "hinge region" situated between the first and second domains of the constant region.

Among the proteins forming the class of the immunoglobulins, most of them are antibodies and accordingly present an antigen binding site or several antigen binding sites.

According to the four-chain model, the antigen binding site of an antibody is located in the variable domains of each of the heavy and light chains, and

requires the association of the heavy and the light chains variable domains.

For the definition of these four-chain model immunoglobulins, reference is made to Roitt. I et al (Immunology-second-Edition Gower Medical Publishing USA, 1989). Reference is especially made to the part concerning the definition of the four-chain immunoglobulins, their polypeptidic and genetic structures, the definition of their variable and constant regions and the obtention of the fragments produced by enzymatic degradation according to well known techniques.

The inventors have surprisingly established that different molecules can be isolated from animals which naturally produce them, which molecules have functional properties of immunoglobulins these functions being in some cases related to structural elements which are distinct from those involved in the function of four-chain immunoglobulins due for instance to the absence of light chains.

The invention relates to two-chain model immunoglobulins which neither correspond to fragments obtained for instance by the degradation in particular the enzymatic degradation of a natural four-chain model immunoglobulin, nor correspond to the expression in host cells, of DNA coding for the constant or the variable region of a natural four-chain model immunoglobulin or a part of these regions, nor correspond to antibodies produced in lymphopaties for example in mice, rats or human.

E.S. Ward et al (1) have described some experiments performed on variable domains of heavy polypeptide chains (V_H) or/and light polypeptide chains (V_L/F_L) to test the ability of these variable domains, to bind specific antigens. For this purpose, a library of V_H genes was prepared from the spleen

genomic DNA of mice previously immunized with these specific antigens.

Ward et al have described in their publication that V_H domains are relatively sticky, presumably due to the exposed hydrophobic surface normally capped by the V_K or V_L domains. They consequently envisage that it should be possible to design V_H domains having improved properties and further that V_H domains with binding activities could serve as the building blocks for making variable fragments (Fv fragments) or complete antibodies.

The invention does not start from the idea that the different fragments (light and heavy chains) and the different domains of these fragments of four-chain model immunoglobulin can be modified to define new or improved antigen binding sites or a four-chain model immunoglobulin.

The inventors have determined that immunoglobulins can have a different structure than the known four-chain model and that such different immunoglobulins offer new means for the preparation of diagnosis reagents, therapeutical agents or any other reagent for use in research or industrial purposes.

Thus the invention provides new immunoglobulins which are capable of showing functional properties of four-chain model immunoglobulins although their structure appears to be more appropriate in many circumstances for their use, their preparation and in some cases for their modification. Moreover these molecules can be considered as lead structures for the modification of other immunoglobulins. The advantages which are provided by these immunoglobulins comprise the possibility to prepare them with an increased facility.

The invention accordingly relates to immunoglobulins characterized in that they comprise two heavy polypeptide chains sufficient for the

formation of a complete antigen binding site or several antigen binding sites, these immunoglobulins being further devoid of light polypeptide chains. In a particular embodiment of the invention, these immunoglobulins are further characterized by the fact that they are the product of the expression in a prokaryotic or in a eukaryotic host cell, of a DNA or of a cDNA having the sequence of an immunoglobulin devoid of light chains as obtainable from lymphocytes or other cells of Camelids.

The immunoglobulins of the invention can be obtained for example from the sequences which are described in figure 7.

The immunoglobulins of the invention, which are devoid of light chains are such that the variable domains of their heavy chains have properties differing from those of the four-chain immunoglobulin V_H . The variable domain of a heavy-chain immunoglobulin of the invention has no normal interaction sites with the V_L or with the C_H1 domain which do not exist in the heavy chain immunoglobulins. it is hence a novel fragment in many of its properties such as solubility and position of the binding site. For clarity reasons we will call it V_{HH} in this text to distinguish it from the classical V_H of four-chain immunoglobulins.

By "a complete antigen binding site" it is meant according to the invention, a site which will alone allow the recognition and complete binding of an antigen. This could be verified by any known method regarding the testing of the binding affinity.

These immunoglobulins which can be prepared by the technique of recombinant DNA, or isolated from animals, will be sometimes called "heavy-chain immunoglobulins" in the following pages. In a preferred embodiment of the invention, these immunoglobulins are in a pure form.

In a first embodiment, the immunoglobulins of the invention are obtainable in prokaryotic cells, especially in E.coli cells by a process comprising the steps of :

- a) cloning in a Bluecript vector of a DNA or cDNA sequence coding for the V_{HH} domain of an immunoglobulin devoid of light chain obtainable for instance from lymphocytes of Camelids,
- b) recovering the cloned fragment after amplification using a 5' primer containing an Xho site and a 3' primer containing the Spe site having the following sequence
TC TTA ACT AGT GAG GAG ACG GTG ACC TG,
- c) cloning the recovered fragment in phase in the immuno PBS vector after digestion of the vector with Xho and Spe restriction enzymes,
- d) transforming host cells, especially E.coli by transfection with the recombinant immuno PBS vector of step c,
- e) recovering the expression product of the V_{HH} coding sequence, for instance by using antibodies raised against the dromadary V_{HH} domain.

In another embodiment the immunoglobulins are hetero-specific immunoglobulins obtainable by a process comprising the steps of:

- obtaining a first DNA or cDNA sequence coding for a V_{HH} domain or part thereof having a determined specificity against a given antigen and comprised between Xho and Spe sites,
- obtaining a second DNA or cDNA sequence coding for a V_{HH} domain or part thereof, having a determined specificity different from the specificity of the first DNA or cDNA sequence and comprised between the Spe and EcoRI sites,
- digesting an immuno PBS vector with EcoRI and XhoI restriction enzymes,

- ligating the obtained DNA or cDNA sequences coding for V_{HH} domains, so that the DNA or cDNA sequences are serially cloned in the vector,
- transforming a host cell, especially E.coli cell by transfection, and recovering the obtained immunoglobulins.

In another embodiment, the immunoglobulins are obtainable by a process comprising the steps of:

- obtaining a DNA or cDNA sequence coding for a V_{HH} domain or part thereof, having a determined specific antigen binding site,
- amplifying the obtained DNA or cDNA, using a 5' primer containing an initiation codon and a HindIII site, and a 3' primer containing a termination codon having a XhoI site,
- recombining the amplified DNA or cDNA into the HindIII (position 2650) and XhoI (position 4067) sites of a plasmid pMM984,
- transfecting permissive cells especially NB-E cells with the recombinant plasmid,
- recovering the obtained products.

Successful expression can be verified with antibodies directed against a region of a V_{HH} domain, especially by an ELISA assay.

According to another particular embodiment of this process, the immunoglobulins are cloned in a parvovirus.

In another example these immunoglobulins are obtainable by a process comprising the further cloning of a second DNA or cDNA sequence having another determined antigen binding site, in the pMM984 plasmid.

Such an Immunoglobulin can be further characterized in that it is obtainable by a process wherein the vector is Yep 52 and the transformed recombinant cell is a yeast especially S.cerevisiae.

A particular Immunoglobulin is characterized in that it has a catalytic activity, especially in that it is directed against an antigen mimicking an activated state of a given substrate. These catalytic antibodies can be modified at the level of their binding site, by random or directed mutagenesis in order to increase or modify their catalytic function. Reference may be made to the publication of Lerner et al (TIBS November 1987. 427-430) for the general technique for the preparation of such catalytic immunoglobulins.

According to a preferred embodiment, the immunoglobulins of the invention are characterized in that their variable regions contain in position 45, an amino-acid which is different from leucine, proline or glutamine residue.

Moreover the heavy-chain immunoglobulins are not products characteristic of lymphocytes of animals nor from lymphocytes of a human patient suffering from lymphopathies. Such immunoglobulins produced in lymphopathies are monoclonal in origin and result from pathogenic mutations at the genomic level. They have apparently no antigen binding site.

The two heavy polypeptide chains of these immunoglobulins can be linked by a hinge region according to the definition of Roitt et al.

In a particular embodiment of the invention, immunoglobulins corresponding to the above-defined molecules are capable of acting as antibodies.

The antigen binding site(s) of the immunoglobulins of the invention are located in the variable region of the heavy chain.

In a particular group of these immunoglobulins each heavy polypeptide chain contains one antigen binding site on its variable region, and these sites correspond to the same amino-acid sequence.

In a further embodiment of the invention the immunoglobulins are characterized in that their heavy polypeptide chains contain a variable region (V_{HH}) and a constant region (C_H) according to the definition of Roitt et al, but are devoid of the first domain of their constant region. This first domain of the constant region is called C_{H1} .

These immunoglobulins having no C_{H1} domain are such that the variable region of their chains is directly linked to the hinge region at the C-terminal part of the variable region.

The immunoglobulins of the type described hereabove can comprise type G immunoglobulins and especially immunoglobulins which are defined as immunoglobulins of class 2 (IgG2) or immunoglobulins of class 3 (IgG3).

The absence of the light chain and of the first constant domain lead to a modification of the nomenclature of the immunoglobulin fragments obtained by enzymatic digestion, according to Roitt et al.

The terms Fc and pFc on the one hand, Fc' and pFc' on the other hand corresponding respectively to the papain and pepsin digestion fragments are maintained.

The terms Fab $F(ab)_2$ $F(ab')_2$ Fabc, Fd and Fv are no longer applicable in their original sense as these fragments have either a light chain, the variable part of the light chain or the C_{H1} domain.

The fragments obtained by papain digestion and composed of the V_{HH} domain and the hinge region will be called $FV_{HH}h$ or $F(V_{HH}h)_2$ depending upon whether or not they remain linked by the disulphide bonds.

In another embodiment of the invention, immunoglobulins replying to the hereabove given definitions can be originating from animals especially from animals of the camelid family. The inventors have found out that the heavy-chain immunoglobulins which

are present in camelids are not associated with a pathological situation which would induce the production of abnormal antibodies with respect to the four-chain immunoglobulins. On the basis of a comparative study of old world camelids (Camelus bactrianus and Camelus dromaderius) and new world camelids (for example Lama Paccos, Lama Glama, and Lama Vicugna) the inventors have shown that the immunoglobulins of the invention, which are devoid of light polypeptide chains are found in all species. Nevertheless differences may be apparent in molecular weight of these immunoglobulins depending on the animals. Especially the molecular weight of a heavy chain contained in these immunoglobulins can be from approximately 43 kd to approximately 47 kd, in particular 45 kd.

Advantageously the heavy-chain immunoglobulins of the invention are secreted in blood of camelids.

Immunoglobulins according to this particular embodiment of the invention are obtainable by purification from serum of camelids and a process for the purification is described in details in the examples. In the case where the immunoglobulins are obtained from Camelids, the invention relates to immunoglobulins which are not in their natural biological environment.

According to the invention immunoglobulin IgG2 as obtainable by purification from the serum of camelids can be characterized in that :

- it is not adsorbed by chromatography on Protein G Sepharose column,
- it is adsorbed by chromatography on Protein A Sepharose column,
- it has a molecular weight of around 100 kd after elution with a pH 4.5 buffer (0.15 M NaCl, 0.58% acetic acid adjusted to pH 4.5 by NaOH),

- it consists of heavy $\gamma 2$ polypeptide chains of a molecular weight of around 46 kd preferably 45 after reduction.

According to a further embodiment of the invention another group of immunoglobulins corresponding to IgG3, as obtainable by purification from the serum of Camelids is characterized in that the immunoglobulin :

- is adsorbed by chromatography on a Protein A Sepharose column,
- has a molecular weight of around 100 kd after elution with a pH 3.5 buffer (0.15 M NaCl, 0.58% acetic acid),
- is adsorbed by chromatography on a Protein G Sepharose column and eluted with pH 3.5 buffer (0.15 M NaCl, 0.58% acetic acid).
- consists of heavy $\gamma 3$ polypeptide chains of a molecular weight of around 45 Kd in particular between 43 and 47 kd after reduction.

The immunoglobulins of the invention which are devoid of light chains, nevertheless comprise on their heavy chains a constant region and a variable region. The constant region comprises different domains.

The variable region of immunoglobulins of the invention comprises frameworks (FW) and complementarity determining regions (CDR), especially 4 frameworks and 3 complementarity regions. It is distinguished from the four-chain immunoglobulins especially by the fact that this variable region can itself contain an antigen binding site or several, without contribution of the variable region of a light chain which is absent.

The amino-acid sequences of frameworks 1 and 4 comprise among others respectively amino-acid sequences which can be selected from the following :

for the framework 1 domain

11

GGSVQTGGSLRLSCEISGLTFD
GGSVQTGGSLRLSCAVSGFSFS
GGSEQGGGSLRLSCAISGYTYG
GGSVQPGGSLTSLCTVSGATYS
GGSVQAGGSLRLSCTGSGFPYS
GGSVQAGGSLRLSCVAGFGTS
GGSVQAGGSLRLSCVSPSS

for the framework 4 domain

WGQGTQVTVSS
WGQGTLVTVSS
WGQGAQVTVSS
WGQGTQVTASS
RGQGTQVTVSL

for the CDR3 domain

ALQPGGYCGYGX - - - - - CL
VSLMDRISQH - - - - - GC
VPAHLGPGAILD LKKY - - - - - KY
FCYSTAGDGGSGE - - - - - MY
ELSGGSC ELP L LF - - - - - DY
DWKYWTCGAQTGGYF - - - - - GQ
RLTEMGACDARWATLATRTFAYNY
QKKDRTRWAEPREW - - - - - NN
GSRFSSPVGSTSRLES - SDY - - NY
ADPSIYY S I L X I E Y - - - - - KY
DSPCYMPTMPAPPPIRDSFGW - - DD
TSSFYWYCTTAPY - - - - - NV
TEIEWYGCNLR TTF - - - - - TR
NQLAGGWYLDPNYWLSVGAY - - AI
RLTEMGACDARWATLATRTFAYNY
DGWTRKEGGIGLPWSVQCEDGYNY
DSYPCHLL - - - - - DV
VEYPIADMCS - - - - - RY

As stated above, the immunoglobulins of the invention are preferably devoid of the totality of their C_H1 domain.

Such immunoglobulins comprise C_H2 and C_H3 domains in the C-terminal region with respect to the hinge region.

According to a particular embodiment of the invention the constant region of the immunoglobulins comprises C_H2 and C_H3 domains comprising an amino-acid sequence selected from the following :

for the C_H2 domain:

APELLGGPTVFIFPPKPKDVLSITLTP

APELPGGPSVFVFPTKPKDVLSISGRP

APELPGGPSVFVFPPKPKDVLSISGRP

APELLGGPSVFIFPPKPKDVLSISGRP

for the C_H3 domain:

GQTREPQVYTLA

GQTREPQVYTLAPXRLEL

GQPREPQVYTLPPSRDEL

GQPREPQVYTLPPSREEM

GQPREPQVYTLPPSQEEM

Interestingly the inventors have shown that the hinge region of the immunoglobulins of the invention can present variable lengths. When these immunoglobulins act as antibodies, the length of the hinge region will participate to the determination of the distance separating the antigen binding sites.

Preferably an immunoglobulin according to the invention is characterized in that its hinge region comprises from 0 to 50 amino-acids.

Particular sequences of hinge region of the immunoglobulins of the invention are the following.

GTNEVCKCPKCP

or,

EPKIPQPQPKPQPQPQPKPQPKPEPECTCPKCP

The short hinge region corresponds to an IgG3 molecule and the long hinge sequence corresponds to an IgG2 molecule.

Isolated V_{HH} derived from heavy chain immunoglobulins or V_{HH} libraries corresponding to the heavy chain immunoglobulins can be distinguished from V_{HH} cloning of four-chain model immunoglobulins on the basis of sequence features characterizing heavy chain immunoglobulins.

The camel heavy - chain immunoglobulin V_{HH} region shows a number of differences with the V_{HH} regions derived from 4-chain immunoglobulins from all species examined. At the levels of the residues involved in the V_{HH}/V_L interactions, an important difference is noted at the level of position 45 (FW) which is practically always leucine in the 4-chain immunoglobulins (98%), the other amino acids at this position being proline (1%) or glutamine (1%).

In the camel heavy-chain immunoglobulin, in the sequences examined at present, leucine at position 45 is only found once. It could originate from a four-chain immunoglobulin. In the other cases, it is replaced by arginine, cysteine or glutamic acid residue. The presence of charged amino acids at this position should contribute to making the V_{HH} more soluble.

The replacement by camelid specific residues such as those of position 45 appears to be interesting for the construction of engineered V_{HH} regions derived from the V_{HH} repertoire of 4-chain immunoglobulins.

A second feature specific of the camelid V_{HH} domain is the frequent presence of a cysteine in the CDR₃ region associated with a cysteine in the CDR₁ position 31 or 33 or FW₂ region at position 45. The possibility of establishing a disulphide bond between the CDR₃ region and the rest of the variable domain

would contribute to the stability and positioning of the binding site.

With the exception of a single pathogenic myeloma protein (DAW) such a disulphide bond has never been encountered in immunoglobulin V regions derived from 4 chain immunoglobulins.

The heavy-chain immunoglobulins of the invention have further the particular advantage of being not sticky. Accordingly these immunoglobulins being present in the serum, aggregate much less than isolated heavy chains of a four-chain immunoglobulins. The immunoglobulins of the invention are soluble to a concentration above 0.5 mg/ml, preferably above 1 mg/ml and more advantageously above 2 mg/ml.

These immunoglobulins further bear an extensive antigen binding repertoire and undergo affinity and specificity maturation in vivo. Accordingly they allow the isolation and the preparation of antibodies having defined specificity, regarding determined antigens.

Another interesting property of the immunoglobulins of the invention is that they can be modified and especially humanized. Especially it is possible to replace all or part of the constant region of these immunoglobulins by all or part of a constant region of a human antibody. For example the C_H2 and/or C_H3 domains of the immunoglobulin could be replaced by the C_H2 and/or C_H3 domains of the IgG γ 3 human immunoglobulin.

In such humanized antibodies it is also possible to replace a part of the variable sequence, namely one or more of the framework residues which do not intervene in the binding site by human framework residues, or by a part of a human antibody.

Conversely features (especially peptide fragments) of heavy-chain immunoglobulin V_{HH} regions, could be introduced into the V_H or V_L regions derived from four-chain immunoglobulins with for instance the

aim of achieving greater solubility of the immunoglobulins.

The invention further relates to a fragment of an immunoglobulin which has been described hereabove and especially to a fragment selected from the following group :

- a fragment corresponding to one heavy polypeptide chain of an immunoglobulin devoid of light chains,
- fragments obtained by enzymatic digestion of the immunoglobulins of the invention, especially those obtained by partial digestion with papain leading to the Fc fragment (constant fragment) and leading to FV_{HH}h fragment (containing the antigen binding sites of the heavy chains) or its dimer F(V_{HH}h)₂, or a fragment obtained by further digestion with papain of the Fc fragment, leading to the pFc fragment corresponding to the C-terminal part of the Fc fragment,
- homologous fragments obtained with other proteolytic enzymes,
- a fragment of at least 10 preferably 20 amino acids of the variable region of the immunoglobulin, or the complete variable region, especially a fragment corresponding to the isolated V_{HH} domains or to the V_{HH} dimers linked to the hinge disulphide,
- a fragment corresponding to the hinge region of the immunoglobulin, or to at least 6 amino acids of this hinge region,
- a fragment of the hinge region comprising a repeated sequence of Pro-X,
- a fragment corresponding to at least 10 preferably 20 amino acids of the constant region or to the complete constant region of the immunoglobulin.

The invention also relates to a fragment comprising a repeated sequence, Pro-X which repeated sequence contains at least 3 repeats of Pro-X, X being any amino-acid and preferably Gln (glutamine), Lys (lysine) or Glu (acide glutamique); a particular repeated fragment is composed of a 12-fold repeat of the sequence Pro-X.

Such a fragment can be advantageously used as a link between different types of molecules.

The amino-acids of the Pro-X sequence are chosen among any natural or non natural amino-acids.

The fragments can be obtained by enzymatic degradation of the immunoglobulins. They can also be obtained by expression in cells or organisms, of nucleotide sequence coding for the immunoglobulins, or they can be chemically synthesized.

The invention also relates to anti-idiotypes antibodies belonging to the heavy chain immunoglobulin classes. Such anti-idiotypes can be produced against human or animal idiotypes. A property of these anti-idiotypes is that they can be used as idiotypic vaccines, in particular for vaccination against glycoproteins or glycolipids and where the carbohydrate determines the epitope.

The invention also relates to anti-idiotypes capable of recognizing idiotypes of heavy-chain immunoglobulins.

Such anti-idiotypic antibodies can be either syngeneic antibodies or allogenic or xenogeneic antibodies.

The invention also concerns nucleotide sequences coding for all or part of a protein which amino-acid sequence comprises a peptide sequence selected from the following :

G G S V Q T G G S L R L S C E I S G L T F D
G G S V Q T G G S L R L S C A V S G F S F S

GGSEQGGGSLRLSCAISGYTYG
GGSVQPGGSLTSLCTVSGATYS
GGSVQAGGSLRLSCTGSGFPYS
GGSVQAGGSLRLSCVAGFGTS
GGSVQAGGSLRLSCVSFSFSS

WGQGTQVTVSS
WGQGTLVTVSS
WGQGAQVTVSS
WGQGTQVTAASS
RGQGTQVTVSL

ALQPGGYCGYGX-----CL
VSLMDRISQH-----GC
VPAHLGPGA ILDLKKY-----KY
FCYSTAGDGGSGE-----MY
ELSGGSC ELP L LF-----DY
DWKYWTCGAQTGGYF-----GQ
RLTEMGACDARWATLATRTFAYNY
QKKDRTRWAEPREW-----NN
GSRFSSSPVGSTSRLES-SDY--NY
ADPSIYYYSILXIEY-----KY
DSPCYMPTMPAPP I RDSFGW--DD
TSSSFYWYCTTAPY-----NV
TEIEWYGCNLR TTF-----TR
NQLAGGWYLDPNYWLSVGAY--AI
RLTEMGACDARWATLATRTFAYNY
DGWTRKEGGIGLPWSVQCEDGYNY
DSYPCHLL-----DV
VEYP IADMCS-----RY

APELLGGPSVFVFPKPKDVLSISGXPK

APELPGGPSVFVFPKPKDVLSISGRPK

APELPGGPSVFVFPKPKDVLSISGRPK

APELLGGPSVFIFPPKPKDVLSISGRPK

GQTRPQVYTLAPXRLEL

GQPREPQVYTLPPSRDEL

GQPREPQVYTLPPSREEM

GQPREPQVYTLPPSQEEM

VTVSSGTNEVCKCPKCPAPELPGGPSVFVFP

or,

VTVSSEPKIPQPKPQPKPQPKPQPKPQPKPEPECTCPKCPAPELLGGPSVFIFP

GTNEVCKCPKCP

APELPGGPSVFVFP

EPKIPQPKPQPKPQPKPQPKPQPKPEPECTCPKCP

APELLGGPSVFIFP

Such nucleotide sequences can be deduced from the amino-acid sequences taking into account the denegeracy of the genetic code. They can be synthesized or isolated from cells producing immunoglobulins of the invention.

A procedure for the obtention of such DNA sequences is described in the examples.

The invention also contemplates RNA, especially mRNA sequences corresponding to these DNA sequences, and also corresponding cDNA sequences.

The nucleotide sequences of the invention can further be used for the preparation of primers appropriate for the detection in cells or screening of DNA or cDNA libraries to isolate nucleotide sequences coding for immunoglobulins of the invention.

Such nucleotide sequences can be used for the preparation of recombinant vectors and the expression of these sequences contained in the vectors by host cells especially prokaryotic cells like bacteria or also eukaryotic cells and for example CHO cells, insect cells, simian cells like Vero cells, or any other mammalian cells. Especially the fact that the immunoglobulins of the invention are devoid of light chains permits to secrete them in eukaryotic cells since there is no need to have recourse to the step consisting in the formation of the BIP protein which is required in the four-chain immunoglobulins.

The inadequacies of the known methods for producing monoclonal antibodies or immunoglobulins by recombinant DNA technology comes from the necessity in the vast majority of cases to clone simultaneously the V_H and V_L domains corresponding to the specific binding site of 4 chain immunoglobulins. The animals and especially camelids which produce heavy-chain immunoglobulins according to the invention, and possibly other vertebrate species are capable of producing heavy-chain immunoglobulins of which the binding site is located exclusively in the V_{HH} domain. Unlike the few heavy-chain immunoglobulins produced in other species by chain separation or by direct cloning, the camelid heavy-chain immunoglobulins have undergone extensive maturation in vivo. Moreover their V region has naturally evolved to function in absence of the V_L . They are therefore ideal for producing monoclonal antibodies by recombinant DNA technology. As the obtention of specific antigen binding clones does not depend on a stochastic process necessitating a very large number of recombinant cells, this allows also a much more extensive examination of the repertoire.

This can be done at the level of the non rearranged V_{HH} repertoire using DNA derived from an arbitrarily chosen tissue or cell type or at the level of the rearranged V_{HH} repertoire, using DNA obtained from B lymphocytes. More interesting however is to transcribe the mRNA from antibody producing cells and to clone the cDNA with or without prior amplification into an adequate vector. This will result in the obtention of antibodies which have already undergone affinity maturation.

The examination of a large repertoire should prove to be particularly useful in the search for antibodies with catalytic activities.

The invention thus provides libraries which can be generated in a way which includes part of the hinge sequence, the identification is simple as the hinge is directly attached to the V_{HH} domain.

These libraries can be obtained by cloning cDNA from lymphoid cells with or without prior PCR amplification. The PCR primers are located in the promoter, leader or framework sequences of the V_{HH} for the 5' primer and in the hinge, CH_2 , CH_3 , 3' untranslated region or polyA tail for the 3' primer. A size selection of amplified material allows the construction of a library limited to heavy chain immunoglobulins.

In a particular example, the following 3' primer in which a KpnI site has been constructed and which corresponds to amino-acids 313 to 319 (CGC CAT CAA GGT AAC AGT TGA) is used in conjunction with mouse V_{HH} primers described by Sestry et al and containing a Xho site

AG GTC CAG CTG CTC GAG TCT GG

AG CTC CAG CTG CTC GAG TCT GG

AG GTC CAG CTT CTC GAG TCT GG

XhoI site

These primers yield a library of camelid heavy chain immunoglobulins comprising the V_{HH} region (related to mouse or human subgroup III), the hinge and a section of CH_2 .

In another example, the cDNA is polyadenylated at its 5' end and the mouse specific V_{HH} primers are replaced by a poly T primer with an inbuilt XhoI site, at the level of nucleotide 12.

CTCGAGT₁₂.

The same 3' primer with a KpnI site is used.

This method generates a library containing all subgroups of immunoglobulins.

Part of the interest in cloning a region encompassing the hinge- CH_2 link is that in both $\gamma 2$ and

$\gamma 3$, a Sac site is present immediately after the hinge. This site allows the grafting of the sequence coding for the V_{HH} and the hinge onto the Fc region of other immunoglobulins, in particular the human IgG₁ and IgG₃ which have the same amino acid sequence at this site (Glu₂₄₆ Leu₂₄₇).

As an example, the invention contemplates a cDNA library composed of nucleotide sequences coding for a heavy-chain immunoglobulin, such as obtained by performing the following steps:

a) treating a sample containing lymphoid cells, especially periferal, lymphocytes, spleen cells, lymph nodes or another lymphoid tissue from a healthy animal, especially selected among the Camelids, in order to separate the lymphoid cells,

b) separating polyadenylated RNA from the other nucleic acids and components of the cells,

c) reacting the obtained RNA with a reverse transcriptase in order to obtain the corresponding cDNA,

d) contacting the cDNA of step c) with 5' primers corresponding to mouse V_H domain of four-chain immunoglobulins, which primer contains a determined restriction site, for example an XhoI site and with 3' primers corresponding to the N-terminal part of a C_H2 domain containing a KpnI site,

e) amplifying the DNA,

f) cloning the amplified sequence in a vector, especially in a bluescript vector,

g) recovering the clones hybridizing with a probe corresponding to the sequence coding for a constant domain from an isolated heavy-chain immunoglobulin.

This cloning gives rise to clones containing DNA sequences including the sequence coding for the hinge. It thus permits the characterization of the subclass of the immunoglobulin and the SacI site useful for grafting the FV_{HH} to the Fc region.

The recovery of the sequences coding for the heavy-chain immunoglobulins can also be achieved by the selection of clones containing DNA sequences having a size compatible with the lack of the C_H1 domain.

It is possible according to another embodiment of the invention, to add the following steps between steps c) and d) of the above process:

- in the presence of a DNA polymerase and of deoxyribonucleotide triphosphates, contacting said cDNA with oligonucleotide degenerated primers, which sequences are capable of coding for the hinge region and N-terminal V_{HH} domain of an immunoglobulin, the primers being capable of hybridizing with the cDNA and capable of initiating the extension of a DNA sequence complementary to the cDNA used as template,
- recovering the amplified DNA.

The clones can be expressed in several types of expression vectors. As an example using a commercially available vector Immuno PBS (Huse et al : Science (1989) 246, 1275), clones produced in Bluescript[®] according to the above described procedure, are recovered by PCR using the same XhoI containing 5' primer and a new 3' primer, corresponding to residues 113-103 in the framework of the immunoglobulins, in which an Spe site has been constructed : TC TTA ACT AGT GAG GAG ACG GTG ACC TG. This procedure allows the cloning of the V_{HH} in the Xho/Spe site of the Immuno PBS vector. However, the 3' end of the gene is not in phase with the identification "tag" and the stop codon of the vector. To achieve this, the construct is cut with Spe and the 4 base overhangs are filled in, using the Klenow fragment after which the vector is religated. A further refinement consists in replacing the marker ("tag") with a poly histidine so that metal purification of the cloned V_{HH} can be performed. To

achieve this a Spe/EcoRI double stranded oligo-nucleotide coding for 6 histidines and a termination codon is first constructed by synthesis of both strands followed by heating and annealing :

CTA GTG CAC CAC CAT CAC CAT CAC TAA* TAG*

AC GTG GTG GTA GTG GTA GTG ATT ATC TTA A

The vector containing the insert is then digested with SpeI and EcoRI to remove the resident "tag" sequence which can be replaced by the poly-His/termination sequence. The produced V_{HH} can equally be detected by using antibodies raised against the dromedary V_{HH} regions. Under laboratory conditions, V_{HH} regions are produced in the Immuno PBS vector in mg amounts per liter.

The invention also relates to a DNA library composed of nucleotide sequences coding for a heavy-chain immunoglobulin, such as obtained from cells with rearranged immunoglobulin genes.

In a preferred embodiment of the invention, the library is prepared from cells from an animal previously immunized against a determined antigen. This allows the selection of antibodies having a preselected specificity for the antigen used for immunization.

In another embodiment of the invention, the amplification of the cDNA is not performed prior to the cloning of the cDNA.

The heavy-chain of the four-chain immunoglobulins remains sequestered in the cell by a chaperon protein (BIP) until it has combined with a light chain. The binding site for the chaperon protein is the C_{H1} domain. As this domain is absent from the heavy chain immunoglobulins, their secretion is independent of the presence of the BIP protein or of the light chain. Moreover the inventors have shown that the obtained immunoglobulins are not sticky and accordingly will not abnormally aggregate.

The invention also relates to a process for the preparation of a monoclonal antibody directed against a determined antigen, the antigen binding site of the antibody consisting of heavy polypeptide chains and which antibody is further devoid of light polypeptide chains, which process comprises :

- immortalizing lymphocytes, obtained for example from the peripheral blood of Camelids previously immunized with a determined antigen, with an immortal cell and preferably with myeloma cells, in order to form a hybridoma,
- culturing the immortalized cells (hybridoma) formed and recovering the cells producing the antibodies having the desired specificity.

The preparation of antibodies can also be performed without a previous immunization of Camelids.

According to another process for the preparation of antibodies, the recourse to the technique of the hybridoma cell is not required.

According to such process, antibodies are prepared in vitro and they can be obtained by a process comprising the steps of :

- cloning into vectors, especially into phages and more particularly filamentous bacteriophages, DNA or cDNA sequences obtained from lymphocytes especially PBLs of Camelids previously immunized with determined antigens,
- transforming prokaryotic cells with the above vectors in conditions allowing the production of the antibodies,
- selecting the antibodies for their heavy-chain structure and further by subjecting them to antigen-affinity selection,
- recovering the antibodies having the desired specificity,

In another embodiment of the invention the cloning is performed in vectors, especially into

plasmids coding for bacterial membrane proteins. Prokaryotic cells are then transformed with the above vectors in conditions allowing the expression of antibodies in their membrane.

The positive cells are further selected by antigen affinity selection.

The heavy chain antibodies which do not contain the C_H1 domain present a distinct advantage in this respect. Indeed, the C_H1 domain binds to BIP type chaperone proteins present within eukaryotic vectors and the heavy chains are not transported out of the endocytosolic reticulum unless light chains are present. This means that in eukaryotic cells, efficient cloning of 4-chain immunoglobulins in non mammalian cells such as yeast cells can depend on the properties of the resident BIP type chaperone and can hence be very difficult to achieve. In this respect the heavy chain antibodies of the invention which lack the CH₁ domain present a distinctive advantage.

In a preferred embodiment of the invention the cloning can be performed in yeast either for the production of antibodies or for the modification of the metabolism of the yeast. As example, Yep 52 vector can be used. This vector has the origin of replication (ORI) 2 μ of the yeast together with a selection marker Leu 2.

The cloned gene is under the control of gal1 promoter and accordingly is inducible by galactose. Moreover, the expression can be repressed by glucose which allows the obtention of very high concentration of cells before the induction.

The cloning between BamHI and SalI sites using the same strategy of production of genes by PCR as the one described above, allows the cloning of camelid immunoglobulin genes in E. coli. As example of metabolic modulation which can be obtained by

antibodies and proposed for the yeast, one can site the cloning of antibodies directed against cyclins, that is proteins involved in the regulation of the cellular cycle of the yeast (TIBS 16 430 J.D. Mc Kinney, N. Heintz 1991). Another example is the introduction by genetic engineering of an antibody directed against CD₂₈, which antibody would be inducible (for instance by gall), within the genome of the yeast. The CD₂₈ is involved at the level of the initiation of cell division, and therefore the expression of antibodies against this molecule would allow an efficient control of multiplication of the cells and the optimization of methods for the production in bioreactors or by means of immobilized cells.

In yet another embodiment of the invention, the cloning vector is a plasmid or a eukaryotic virus vector and the cells to be transformed are eukaryotic cells, especially yeast cells, mammalian cells for example CHO cells or simian cells such as Vero cells, insect cells, plant cells, or protozoan cells.

For more details concerning the procedure to be applied in such a case, reference is made to the publication of Marks et al, J. Mol. Biol. 1991, 222:581-597.

Furthermore, starting from the immunoglobulins of the invention, or from fragments thereof, new immunoglobulins or derivatives can be prepared.

Accordingly immunoglobulins replying to the above given definitions can be prepared against determined antigens. Especially the invention provides monoclonal or polyclonal antibodies devoid of light polypeptide chains or antisera containing such antibodies and directed against determined antigens and for example against antigens of pathological agents such as bacteria, viruses or parasites. As example of antigens or antigenic determinants against which antibodies

could be prepared, one can cite the envelope glycoproteins of viruses or peptides thereof, such as the external envelope glycoprotein of a HIV virus, the surface antigen of the hepatitis B virus.

Immunoglobulins of the invention can also be directed against a protein, hapten, carbohydrate or nucleic acid.

Particular antibodies according to the invention are directed against the galactosyl α -1-3-galactose epitope.

The immunoglobulins of the invention allow further the preparation of combined products such as the combination of the heavy-chain immunoglobulin or a fragment thereof with a toxin, an enzyme, a drug, a hormone.

As example one can prepare the combination of a heavy-chain immunoglobulin bearing an antigen binding site recognizing a myeloma immunoglobulin epitope with the abrin or mistletoe lectin toxin. Such a construct would have its uses in patient specific therapy.

Another advantageous combination is that one can prepare between a heavy-chain immunoglobulins recognizing an insect gut antigen with a toxin specific for insects such as the toxins of the different serotypes of *Bacillus thuringiensis* or *Bacillus sphaericus*. Such a construct cloned into plants can be used to increase the specificity or the host range of existing bacterial toxins.

The invention also proposes antibodies having different specificities on each heavy polypeptide chains. These multifunctional, especially bifunctional antibodies could be prepared by combining two heavy chains of immunoglobulins of the invention or one heavy chain of an immunoglobulin of the invention with a fragment of a four-chain model immunoglobulin.

The invention also provides hetero-specific antibodies which can be used for the targetting of

drugs or any biological substance like hormones. In particular they can be used to selectively target hormones or cytokines to a limited category of cells. Examples are a combination of a murine or human antibody raised against interleukin 2 (IL₂) and a heavy-chain antibody raised against CD₄ cells. This could be used to reactivate CD₄ cells which have lost their IL₂ receptor.

The heavy-chain immunoglobulins of the invention can also be used for the preparation of hetero-specific antibodies. These can be achieved either according to the above described method by reduction of the bridges between the different chains and reoxydation, according to the usual techniques, of two antibodies having different specificities, but it can also be achieved by serial cloning of two antibodies for instance in the Immuno pBS vector.

In such a case, a first gene corresponding to the V_{HH} domain comprised between Xho site and a Spe site is prepared as described above. A second gene is then prepared through an analogous way by using as 5' extremity a primer containing a Spe site, and as 3' extremity a primer containing a termination codon and an EcoRI site. The vector is then digested with EcoRI and XhoI and further both V_{HH} genes are digested respectively by Xho/Spe and by Spe/EcoRI.

After ligation, both immunoglobulin genes are serially cloned. The spacing between both genes can be increased by the introduction of addition codons within the 5' SpeI primer.

In a particular embodiment of the invention, the hinge region of IgG2 immunoglobulins according to the invention is semi-rigid and is thus appropriate for coupling proteins. In such an application proteins or peptides can be linked to various substances, especially to ligands through the hinge region used as

spacer. Advantageously the fragment comprises at least 6 amino acids.

According to the invention it is interesting to use a sequence comprising a repeated sequence Pro-X, X being any amino-acid and preferably Gln, Lys or Glu, especially a fragment composed of at least a 3-fold repeat and preferably of a 12-fold repeat, for coupling proteins to ligand, or for assembling different protein domains.

The hinge region or a fragment thereof can also be used for coupling proteins to ligands or for assembling different protein domains.

Usual techniques for the coupling are appropriate and especially reference may be made to the technique of protein engineering by assembling cloned sequences.

The antibodies according to this invention could be used as reagents for the diagnosis in vitro or by imaging techniques. The immunoglobulins of the invention could be labelled with radio-isotopes, chemical or enzymatic markers or chemiluminescent markers.

As example and especially in the case of detection or observation with the immunoglobulins by imaging techniques, a label like technetium, especially technitium 99 is advantageous. This label can be used for direct labelling by a coupling procedure with the immunoglobulins or fragments thereof or for indirect labelling after a step of preparation of a complex with the technitium.

Other interesting radioactive labels are for instance indium and especially indium 111, or iodine, especially I^{131} , I^{125} and I^{123} .

For the description of these techniques reference is made to the FR patent application published under number 2649488.

In these applications the small size of the V_{HH} fragment is a definitive advantage for penetration into tissue.

The invention also concerns monoclonal antibodies reacting with anti-idiotypes of the above-described antibodies.

The invention also concerns cells or organisms in which heavy-chain immunoglobulins have been cloned. Such cells or organisms can be used for the purpose of producing heavy-chain immunoglobulins having a desired preselected specificity, or corresponding to a particular repertoire. They can also be produced for the purpose of modifying the metabolism of the cell which expresses them. In the case of modification of the metabolism of cells transformed with the sequences coding for heavy-chain immunoglobulins, these produced heavy-chain immunoglobulins are used like antisense DNA. Antisense DNA is usually involved in blocking the expression of certain genes such as for instance the variable surface antigen of trypanosomes or other pathogens. Likewise, the production or the activity of certain proteins or enzymes could be inhibited by expressing antibodies against this protein or enzyme within the same cell.

The invention also relates to a modified 4-chain immunoglobulin or fragments thereof, the V_H regions of which has been partially replaced by specific sequences or amino acids of heavy chain immunoglobulins, especially by sequences of the V_{HH} domain. A particular modified V_H domain of a four-chain immunoglobulin, is characterized in that the leucine, proline or glutamine in position 45 of the V_H regions has been replaced by other amino acids and preferably by arginine, glutamic acid or cysteine.

A further modified V_H or V_L domain of a four-chain immunoglobulin, is characterized by linking of CDR loops together or to FW regions by the

introduction of paired cysteines, the CDR region being selected between the CDR₁ and the CDR₃, the FW region being the FW₂ region, and especially in which one of the cysteines introduced is in position 31, 33 of FR₂ or 45 of CDR₂ and the other in CDR₃.

Especially the introduction of paired cysteines is such that the CDR₃ loop is linked to the FW₂ or CDR₁ domain and more especially the cysteine of the CDR₃ of the V_H is linked to a cysteine in position 31 or 33 of FW₂ or in position 45 of CDR₂.

In another embodiment of the invention, plant cells can be modified by the heavy-chain immunoglobulins according to the invention, in order that they acquire new properties or increased properties.

The heavy-chain immunoglobulins of the invention can be used for gene therapy of cancer for instance by using antibodies directed against proteins present on the tumor cells.

In such a case, the expression of one or two V_{HH} genes can be obtained by using vectors derived from parvo or adeno viruses. The parvo viruses are characterized by the fact that they are devoid of pathogenicity or almost not pathogenic for normal human cells and by the fact that they are capable of easily multiplying in cancer cells (Russel S.J. 1990, Immunol. Today II. 196-200).

The heavy-chain immunoglobulins are for instance cloned within HindIII/XbaI sites of the infectious plasmid of the murine MVM virus (pMM984). (Merchlsinsky et al, 1983, J. Virol. 47, 227-232) and then placed under the control of the MVM38 promoter.

The gene of the V_{HH} domain is amplified by PCR by using a 5' primer containing an initiation codon and a HindIII site, the 3' primer containing a termination codon and a XbaI site.

This construct is then inserted between positions 2650 (HindIII) and 4067 (XbaI) within the plasmid.

The efficiency of the cloning can be checked by transfection. The vector containing the antibody is then introduced in permissive cells (NB-E) by transfection.

The cells are recovered after two days and the presence of V_{HH} regions is determined with an ELISA assay by using rabbit antiserum reacting with the V_{HH} part.

The invention further allows the preparation of catalytic antibodies through different ways. The production of antibodies directed against components mimicking activated states of substrates (as example vanadate as component mimicking the activated state of phosphate in order to produce their phosphoesterase activities, phosphonate as compound mimicking the peptidic binding in order to produce proteases) permits to obtain antibodies having a catalytic function. Another way to obtain such antibodies consists in performing a random mutagenesis in clones of antibodies for example by PCR, in introducing abnormal bases during the amplification of clones. These amplified fragments obtained by PCR are then introduced within an appropriate vector for cloning. Their expression at the surface of the bacteria permits the detection by the substrate of clones having the enzymatic activity. These two approaches can of course be combined. Finally, on the basis of the data available on the structure, for example the data obtained by XRay crystallography or NMR, the modifications can be directed. These modifications can be performed by usual techniques of genetic engineering or by complete synthesis. One advantage of the V_{HH} of the heavy chain immunoglobulins of the invention is the fact that they are sufficiently soluble.

The heavy chain immunoglobulins of the invention can further be produced in plant cells, especially in transgenics plants. As example the heavy chain immunoglobulins can be produced in plants using the pMon530 plasmid (Roger et al. Meth Enzym 153 1566 1987) constitutive plant expression vector as has been described for classical four chain antibodies (Hiat et al. Nature 342 76-78, 1989) once again using the appropriate PCR primers as described above, to generate a DNA fragment in the right phase.

Other advantages and characteristics of the invention will become apparent in the examples and figures which follow.

FIGURES

Figure 1 : Characterisation and purification of camel IgG by affinity chromatography on Protein A and Protein G sepharose (Pharmacia)

(A) shows, after reduction, the SDS-PAGE protein profile of the adsorbed and non adsorbed fractions of Camelus dromedarius serum. The fraction adsorbed on Protein A and eluted with NaCl 0.15 M acetic acid 0.58% show upon reduction (lane c) three heavy chain components of respectively 50, 46 and 43 Kd and light chain (rabbit IgG in lane a). The fractions adsorbed on a Protein G Sepharose (Pharmacia) derivative which has been engineered to delete the albumin binding region (lane e) and eluted with 0.1 M gly HCl pH 2.7 lacks the 46 Kd heavy chain which is recovered in the non adsorbed fraction (lane f). None of these components are present in the fraction non adsorbed on Protein A (lane d), lane b contains the molecular weight markers.

(B) and (C) By differential elution, immunoglobulin fractions containing the 50 and 43 Kd heavy chain can be separated. 5 ml of C. dromedarius serum is adsorbed onto a 5 ml Protein G sepharose column and the column is extensively washed with 20mM phosphate buffer, pH 7.0. Upon elution with pH 3.5 buffer (0.15 M NaCl, 0.58% acetic acid) a 100 Kd component is eluted which upon reduction yields a 43 Kd heavy chain, (lane 1).

After column eluant absorbance has fallen to background level a second immunoglobulin component of 170 Kd can be eluted with pH 2.7 buffer (0.1 M glycine HC). This fraction upon reduction yields a 50 Kd heavy chain and a board light chain band (lane 2).

The fraction non adsorbed on Protein G is then brought on a 5 ml Protein A Sepharose column. After washing

and elution with pH 3.5 buffer (0.15 M NaCl, 0.58% acetic acid) a third immunoglobulin of 100 Kd is obtained which consists solely of 46 Kd heavy chains (lane 3).

Figure 2 : Immunoglobulins of Camelus bactrianus, Lama vicugna, Lama glama and Lama pacos to Protein A (A lanes) and to Protein G (G lanes) analyzed on SDS-PAGE before (A) and after reduction (B)

10 μ l of serum obtained from the different species were added to Eppendorf^R tubes containing 10 mg of Protein A or Protein G sepharose suspended in 400 μ l of pH 8.3 immunoprecipitation buffer (NaCl 0.2. M, Tris 0.01 M; EDTA 0.01 M, Triton X100 1%, ovalbumin 0.1%). The tubes were slowly rotated for 2 hours at 4°C. After centrifugation the pellets were washed 3 times in buffer and once in buffer in which the Triton and ovalbumin had been omitted. The pellets were then resuspended in the SDS-PAGE sample solution 70 μ l per pellet with or without dithiotreitol as reductant. After boiling for 3 min at 100°C, the tubes were centrifuged and the supernatants analysed.

In all species examined the unreduced fractions (A) contain in addition to molecules of approximately 170 Kd also smaller major components of approximately 100 Kd. In the reduced sample (B) the constituent heavy and light chains are detected. In all species a heavy chain component (marked by an asterisk *) is present in the material eluted from the Protein A but absent in the material eluted from the Protein G.

Figure 3 : IgG₁, IgG₂ and IgG₃ were prepared from serum obtained from healthy or Trypanosoma evansi infected Camelus dromedarius (CATT titer 1/160 (3) and analysed by radioimmunoprecipitation

or Western Blotting for anti
trypanosome activity

(A) ^{35}S methionine labelled Trypanosome evansi antigens lysate (500.000 counts) was added to Eppendorf tubes containing 10 μl of serum or, 20 μg of IgG_1 , IgG_2 or IgG_3 in 200 μl of pH 8.3 immunoprecipitation buffer containing 0.1 M TLCK as proteinase inhibitor and slowly rotated at 4°C during one hour. The tubes were then supplemented with 10 mg of Protein A Sepharose suspended in 200 μl of the same pH 8.3 buffer and incubated at 4°C for an additional hour.

After washing and centrifugation at 15000 rpm for 12 s, each pellet was resuspended in 75 μl SDS-PAGE sample solution containing DTT and heated for 3 min. at 100°C. After centrifugation in an Eppendorf minifuge at 15000 rpm for 30 s, 5 μl of the supernatant was saved for radioactivity determination and the remainder analysed by SDS-PAGE and fluorography. The counts/5 μl sample are inscribed on for each line.

(B) 20 μg of IgG_1 , IgG_2 and IgG_3 from healthy and trypanosome infected animals were separated by SDS-PAGE without prior reduction or heating. The separated samples were then electro transferred to a nitrocellulose membrane, one part of the membrane was stained with Ponceau Red to localise the protein material and the remainder incubated with 1% ovalbumin in TST buffer (Tris 10 mM, NaCl 150 mM, Tween 0.05%) to block protein binding sites.

After blocking, the membrane was extensively washed with TST buffer and incubated for 2 hours with ^{35}S -labelled trypanosome antigen. After extensive washing, the membrane was dried and analysed by autoradiography. To avoid background and unspecific binding, the labelled trypanosome lysate was filtered

through a 45 μ millipore filter and incubated with healthy camel immunoglobulin and ovalbumin adsorbed on a nitrocellulose membrane.

figure 4 : Purified IgG3 of the camel, by affinity chromatography on Protein A Sepharose are partially digested with papain and separated on Protein A sepharose.

14mg of purified IgG3 were dissolved in 0.1M phosphate buffer pH 7.0 containing 2mM EDTA. They were digested by 1 hour incubation at 37°C with mercurypapain (1% enzyme to protein ratio) activated by $5 \cdot 10^{-4}$ M cysteine. The digestion was blocked by the addition of excess iodoacetamide ($4 \cdot 10^{-2}$ M) (13). After centrifugation of the digest in an ependorf centrifuge for 5min at 15000 rpm, the papain fragments were separated on a protein A Sepharose column into binding (B) and non binding (NB) fractions. The binding fraction was eluted from the column with 0.1M glycine HCl buffer pH 1.7.

Figure 5 : Schematic presentation of a model for IgG3 molecules devoid of light chains.

Figure 6 : . Schematic representation of immunoglobulins having heavy polypeptide chains and devoid of light chains, regarding conventional four-chain model immunoglobulin
. Representation of a hinge region.

Figure 7: Alignement of 17 V_{HH} DNA sequences of Camel heavy chain immunoglobulins

Figure 8: Expression and purification of the camel V_{HH21} protein from E.coli

I HEAVY CHAIN ANTIBODIES IN CAMELIDS

When Camelus dromedarius serum is adsorbed on Protein G sepharose, an appreciable amount (25-35%) of immunoglobulins (Ig) remains in solution which can then be recovered by affinity chromatography on Protein A sepharose (fig. 1A). The fraction adsorbed on Protein G can be differentially eluted into a tightly bound fraction (25%) consisting of molecules of an unreduced apparent molecular weight (MW) of 170 Kd and a more weakly bound fraction (30-45%) having an apparent molecular weight of 100 Kd (fig. 1B). The 170 Kd component when reduced yields 50 Kd heavy chains and large 30 Kd light chains. The 100 Kd fraction is totally devoid of light chains and appears to be solely composed of heavy chains which after reduction have an apparent MW of 43 Kd (Fig. 1C). The fraction which does not bind to Protein G can be affinity purified and eluted from a Protein A column as a second 100 Kd component which after reduction appears to be composed solely of 46 Kd heavy chains.

The heavy chain immunoglobulins devoid of light chains total up to 75% of the molecules binding to Protein A.

As all three immunoglobulins bind to Protein A we refer to them as IgG : namely IgG₁ (light chain and heavy chain γ 1 (50 Kd) binding to Protein G, IgG₂ (heavy chain γ 2 (46 Kd) non binding to Protein G and IgG₃ (heavy chain γ 3 (43 Kd) binding to Protein G. There is a possibility that these three sub(classes) can be further subdivided.

A comparative study of old world camelids (Camelus bactrianus and Camelus dromedarius) and new world camelids (Lama pacos, Lama glama, Lama vicugna) showed that heavy chain immunoglobulins are found in all species examined, albeit with minor differences in apparent molecular weight and proportion. The new

world camelids differs from the old world camelids in having a larger IgG₃ molecule (heavy chain immunoglobulin binding to Protein G) in which the constituent heavy chains have an apparent molecular weight of 47 Kd (fig. 2).

The abundance of the heavy chain immunoglobulins in the serum of camelids raises the question of what their role is in the immune response and in particular whether they bear antigen binding specificity and if so how extensive is the repertoire. This question could be answered by examining the immunoglobulins from Trypanosoma evansi infected camels (Camelus dromedarius).

For this purpose, the corresponding fractions of IgG₁, IgG₂, IgG₃ were prepared from the serum of a healthy camel and from the serum of camels with a high antitrypanosome titer, measured by the Card Agglutination Test (3). In radio-immunoprecipitation, IgG₁, IgG₂ and IgG₃ derived from infected camel indicating extensive repertoire heterogeneity and complexity (Fig. 3A) were shown to bind a large number of antigens present in a ³⁵S methionine labelled trypanosome lysate.

In blotting experiments ³⁵S methionine labelled trypanosome lysate binds to SDS PAGE separated IgG₁, IgG₂ and IgG₃ obtained from infected animals (Fig. 3B).

This leads us to conclude that the camelid heavy chain IgG₂ and IgG₃ are bona fide antigen binding antibodies.

An immunological paradigm states that an extensive antibody repertoire is generated by the combination of the light and heavy chain variable V region repertoires (6). The heavy chain immunoglobulins of the camel seem to contradict this paradigm.

Immunoglobulins are characterized by a complex I.E.F. (isoelectric focussing) pattern reflecting their extreme heterogeneity. To determine whether the two heavy chains constituting the IgG₂ and IgG₃ are identical or not, the isoelectric focussing (I.E.F.) pattern were observed before and after chain separation by reduction and alkylation using iodoacetamide as alkylating agent.

As this alkylating agent does not introduce additional charges in the molecule, the monomers resulting from the reduction and alkylation of a heavy chain homodimer will have practically the same isoelectric point as the dimer, whereas if they are derived from a heavy chain heterodimer, the monomers will in most cases differ sufficiently in isoelectric point to generate a different pattern in I.E.F.

Upon reduction, and alkylation by iodoacetamide the observed pattern is not modified for the Camelus dromedarius IgG₂ and IgG₃ indicating that these molecules are each composed of two identical heavy chains which migrate to the same position as the unreduced molecule they originated from.

In contrast, the I.E.F. pattern of IgG₁ is completely modified after reduction as the isoelectric point of each molecule is determined by the combination of the isoelectric points of the light and heavy chains which after separation will each migrate to a different position.

These findings indicate that the heavy chains alone can generate an extensive repertoire and question the contribution of the light chain to the useful antibody repertoire. If this necessity be negated, what other role does the light chain play.

Normally, isolated heavy chain from mammalian immunoglobulins tend to aggregate considerably but are only solubilized by light chains (8, 9) which bind to the C_H1 domain of the heavy chain.

In humans and in mice a number of spontaneous or induced myelomas produce a pathological immunoglobulin solely composed of heavy chains (heavy chain disease). These myeloma protein heavy chains carry deletions in the C_H1 and V_{HH} domains (10). The reason why full length heavy chains do not give rise to secreted heavy chain in such pathological immunoglobulins seems to stem from the fact that the synthesis of Ig involves a chaperoning protein, the immunoglobulin heavy chain binding protein or BIP (11), which normally is replaced by the light chain (12). It is possible that the primordial role of the light chain in the four-chain model immunoglobulins is that of a committed heavy chain chaperon and that the emergence of light chain repertoires has just been an evolutionary bonus.

The camelid $\gamma 2$ and $\gamma 3$ chains are considerably shorter than the normal mammalian γ chain. This would suggest that deletions have occurred in the C_H1 domain. Differences in sizes of the $\gamma 2$ and $\gamma 3$ immunoglobulins of old and new world camelids suggests that deletions occurred in several evolutionary steps especially in the C_H1 domain.

II THE HEAVY CHAIN IMMUNOGLOBULINS OF THE CAMELIDS LACK THE C_H1 DOMAIN.

The strategy followed for investigating the heavy chain immunoglobulin primary structure is a combination of protein and cDNA sequencing ; the protein sequencing is necessary to identify sequence stretches characteristic of each immunoglobulin. The N-terminal of the immunoglobulin being derived from the heavy chain variable region repertoire only yields information on the V_{HH} subgroups (variable region of the heavy chain) and cannot be used for class or subclass identification. This means that sequence data

had to be obtained from internal enzymatic or chemical cleavage sites.

A combination of papain digestion and Protein A affinity chromatography allowed the separation of various fragments yielding information on the general structure of IgG3.

The IgG3 of the camel (Camelus dromedarius) purified by affinity chromatography on Protein A Sepharose were partially digested with papain and the digest was separated on Protein A Sepharose into binding and non binding fractions. These fractions were analysed by SDS PAGE under reducing and non reducing conditions (fig 4).

The bound fraction contained two components, one of 28 Kd and one of 14.4 Kd, in addition to uncleaved or partially cleaved material. They were well separated by gel electrophoresis (from preparative 19% SDS-PAGE gels) under non reducing conditions and were further purified by electroelution(in 50mM ammonium bicarbonate, 0.1% (w/v) SDS using a BioRad electro-eluter). After lyophilization of these electroeluted fractions, the remaining SDS was eliminated by precipitating the protein by the addition of 90% ethanol, mixing and incubating the mixture overnight at -20°C (14). The precipitated protein was collected in a pellet by centrifuging (15000 rpm, 5min) and was used for protein sequencing. N-terminal sequencing was performed using the automated Edman chemistry of an Applied Biosystem 477A pulsed liquid protein sequencer. Amino acids were identified as their phenylthiohydantoin (PTH) derivatives using an Applied Biosystem 120 PTH analyser. All chemical and reagents were purchased from Applied Biosystems. Analysis of the chromatographic data was performed using Applied Biosystems software version 1.61. In every case the computer aided sequence analysis was confirmed by

direct inspection of the chromatograms from the PTH analyser. Samples for protein sequencing were dissolved in either 50% (v/v) trifluoroacetic acid (TFA) (28Kd fragment) or 100% TFA (14Kd fragment). Samples of dissolved protein equivalent to 2000 pmol (28Kd fragment) or 500 pmol (14Kd fragment) were applied to TFA-treated glass fibre discs. The glass fibre discs were coated with BioBrene (3mg) and precycled once before use.

N-terminal sequencing of the 28 Kd fragment yields a sequence homologous to the N-terminal part of γ C_H2 domain and hence to the N-terminal end of the Fc fragment. The N-terminal sequence of the 14.4 Kd fragment corresponds to the last lysine of a γ C_H2 and the N-terminal end of a γ C_H3 domain (Table 1). The molecular weight (MW) of the papain fragments and the identification of their N-terminal sequences led us to conclude that the C_H2 and C_H3 domains of the γ 3 heavy chains are normal in size and that the deletion must occur either in the C_H1 or in the V_{HH} domain to generate the shorted γ 3 chain. The fractions which do not bind to Protein A Sepharose contain two bands of 34 and 17 Kd which are more diffuse in SDS PAGE indicating that they originate from the variable N-terminal part of the molecule (fig 4).

Upon reduction, a single diffuse band of 17 Kd is found indicating that the 34 Kd is a disulfide bonded dimer of the 17 Kd component. The 34 Kd fragment apparently contains the hinge and the N-terminal domain V_{HH}.

The protein sequence data can be used to construct degenerate oligonucleotide primers allowing PCR amplification of cDNA or genomic DNA.

It has been shown that the cells from camel spleen imprint cells reacted with rabbit and anti camel immunoglobulin sera and that the spleen was hence a site of synthesis of at least one

immunoglobulin class. cDNA was therefore synthesised from camel spleen mRNA. The conditions for the isolation of RNA were the following: total RNA was isolated from the dromedary spleen by the guanidium isothiocyanate method (15). mRNA was purified with oligo T-paramagnetic beads.

cDNA synthesis is obtained using 1 μ g mRNA template, an oligodT primer and reverse transcriptase (BOEHRINGER MAN). Second strand cDNA is obtained using RNase H and E coli DNA polymerase I according to the condition given by the supplier.

Relevant sequences were amplified by PCR: 5ng of cDNA was amplified by PCR in a 100 μ l reaction mixture (10mM Tris-HCl pH 8.3, 50mM KCl, 15mM MgCl₂, 0.01% (w/v) gelatine, 200 μ M of each dNTP and 25 pmoles of each primer) overlaid with mineral oil (Sigma).

Degenerate primers containing EcoRI and KpnI sites and further cloned into pUC 18. After a round of denaturing and annealing (94°C for 5 min and 54°C for 5 min), 2 units of Taq DNA polymerase were added to the reaction mixture before subjecting it to 35 cycles of amplification: 1 min at 94°C (denature) 1min at 54°C (anneal) , 2 min at 72°C (elongate). To amplify DNA sequences between V_{HH} and C_{H2} domains , (# 72 clones), the PCR was performed in the same conditions with the exception that the annealing temperature was increased to 60°C.

One clone examined (#56/36) had a sequence corresponding to the N-terminal part of a C_{H2} domain identical to the sequence of the 28 Kd fragment. The availability of this sequence data allowed the construction of an exact 3' primer and the cloning of the region between the N-terminal end of the V_{HH} and the C_{H2} domain.

5' primers corresponding to the mouse V_{HH} (16) and containing a XhoI restriction site were used in conjunction with the 3' primer in which a KpnI site

had been inserted and the amplified sequences were cloned into pBluescript^R. Clone #56/36 which displayed two internal HaeIII sites was digested with this enzyme to produce a probe to identify PCR positive clones.

After amplification the PCR products were checked on a 1.2% (w/v) agarose gel. Cleaning up of the PCR products included a phenol-chloroform extraction followed by further purification by HPLC (GEN-PAC FAX column, Waters) and finally by using the MERMAID or GENE CLEAN II kit, BIO 101, Inc) as appropriate. After these purification steps, the amplified cDNA was then digested with EcoRI and KpnI for series #56 clones and with XhoI and KpnI for series #72 clones. A final phenol-chloroform extraction preceded the ligation into pUC 18(series #56 clones) or into pBluescript^R (series #72 clones).

All the clones obtained were smaller than the 860 base pairs to be expected if they possessed a complete V_{HH} and C_{H1} region. Partial sequence data corresponding to the N-terminal of the V_{HH} region reveals that out of 20 clones, 3 were identical and possibly not independent. The sequences obtained resemble the human subgroup III and the murine subgroups IIIa and IIIb (Table 2).

Clones corresponding to two different sets of C_{H2} protein sequences were obtained. A first set of sequences (#72/41) had a N-terminal C_{H2} region identical to the one obtained by protein sequencing of the 28 Kd papain fragments of the $\gamma 3$ heavy chain, a short hinge region containing 3 cysteines and a variable region corresponding to the framework (FR4) residues encoded by the J minigenes adjoining the hinge. The C_{H1} domain is entirely lacking. This cDNA corresponds to the $\gamma 3$ chain (Table 4).

In one closely related sequence (#72/1) the proline in position 259 is replaced by threonine.

The sequence corresponding to the C_H3 and the remaining part of the C_H2 was obtained by PCR of the cDNA using as KpnI primer a poly T in which a KpnI restriction site had been inserted at the 5' end. The total sequence of the γ 3 chain corresponds to a molecular weight (MW) which is in good agreement with the data obtained from SDS PAGE electrophoresis.

The sequence of this γ 3 chain presents similarities with other γ chains except that it lacks the C_H1 domain, the V_{HH} domain being adjacent to the hinge.

One or all three of the cysteines could be probably responsible for holding the two γ 3 chains together.

These results have allowed us to define a model for the IgG3 molecule based on sequence and papain cleavage (fig. 5).

Papain can cleave the molecule on each side of the hinge disulfides and also between C_H2 and C_H3. Under non reducing conditions the V_{HH} domains of IgG3 can be isolated as disulfide linked dimer or as monomer depending on the site of papain cleavage.

A second set of clones #72/29 had a slightly different sequence for the C_H2 and was characterized by a very long hinge immediately preceded by the variable domain. This hinge region has 3 cysteines at its C-terminal end in a sequence homologous to the γ 3 hinge. Such second set of clones could represent the IgG2 subclass. For the constant part of the γ 3 and also for the putative γ 2, most clones are identical showing the γ 2 or γ 3 specific sequences. A few clones such as #72/1 however show minor differences. For instance in the case of clones #72/1 two nucleotide differences are detected.

Several V_{HH} regions cDNA's have now been totally or partially sequenced with the exception of a short stretch at the N-terminal end which is primer derived.

Upon translation the majority shows the characteristic heavy chain Ser₂₁ Cys₂₂ and Tyr₉₀ Tyr₉₁ Cys₉₂ sequences, of the intra V_{HH} region disulfide bridge linking residues 22 and 92. All these clones have a sequence corresponding to the framework 4 (FR4) residues of the variable region immediately preceding the postulated hinge sequence (Table 3). This sequence is generated by the J minigenes and is in the majority of cases similar to the sequence encoded by the human and murine J minigenes. The sequence length between region Cys₉₂ and the C-terminal end of the V_{HH} regions is variable and, in the sequences determined, range from 25 to 37 amino-acids as one might expect from the rearrangements of J and D minigenes varying in length.

Several important questions are raised by the sole existence of these heavy chain immunoglobulins in a non pathological situation. First of all, are they bonafide antibodies ? The heavy chain immunoglobulins obtained from trypanosome infected camels react with a large number of parasite antigens as shown in part I of these examples. This implies that the camelid immune system generates an extensive number of binding sites composed of single V_{HH} domains. This is confirmed by the diversity of the V_{HH} regions of the heavy chain immunoglobulins obtained by PCR.

The second question is "how are they secreted ?". The secretion of immunoglobulin heavy chains composing four-chain model immunoglobulins does not occur under normal conditions. A chaperoning protein, the heavy chain binding protein, or BIP protein, prevents heavy chains from being secreted. It is only when the light chain displaces the BIP protein in the endoplasmatic reticulum that secretion can occur(13).

The heavy chain dimer found in the serum of human or mice with the so-called "heavy chain disease" lack

the C_H1 domains thought to harbour the BIP site (14). In the absence of this domain the BIP protein can no longer bind and prevent the transport of the heavy chains.

The presence in camels of a IgG1 class composed of heavy and light chains making up between 25% and 50% of the total IgG molecules also raises the problem as to how maturation and class switching occurs and what the role of the light chain is. The camelid light chain appears unusually large and heterogeneous when examined in SDS PAGE.

The largest dimension of an isolated domain is 40 Å and the maximum attainable span between binding sites of a conventional IgG with C_H1 and V_{HH} will be of the order of 160 Å ($2V_{HH} + 2C_H1$) (19). The deletion of C_H1 domain in the two types of heavy chain antibodies devoid of light chains, already sequenced has, as a result, a modification of this maximum span (fig. 6). In the IgG3 the extreme distance between the extremities of the V_{HH} regions will be of the order of 80 Å ($2V_{HH}$). This could be a severe limitation for agglutinating or cross linking. In the IgG2 this is compensated by the extremely long stretch of hinge, composed of a 12-fold repeat of the sequence Pro-X (where X is Gln, Lys or Glu) and located N-terminal to the hinge disulfide bridges. In contrast, in the human IgG3, the very long hinge which also apparently arose as the result of sequence duplication does not contribute to increase the distance spanning the two binding sites as this hinge is inter-spersed with disulfide bridges.

The single V_{HH} domain could also probably allow considerably rotational freedom of the binding site versus the Fc domain.

Unlike myeloma heavy chains which result probably from C_H1 deletion in a single antibody producing cell, or heavy chain antibodies produced by expression

cloning(15); the camelid heavy chain antibodies (devoid of light chains) have emerged in a normal immunological environment and it is expected that they will have undergone the selective refinement in specificity and affinity accompanying B cell maturation.

Expression and purification of the camel V_{HH}21 (DR21 on figure 7) protein from E.coli

The clones can be expressed in several types of expression vectors. As an example using a commercially available vector Immuno PBS (Huse et al : Science (1989) 246, 1275), clones produced in Bluescript ® according to the above described procedure, have been recovered by PCR using the same XhoI containing 5' primer and a new 3' primer, corresponding to residues 113-103 in the framework of the immunoglobulins, in which an Spe site has been constructed : TC TTA ACT AGT GAG GAG ACG GTG ACC TG. This procedure allowed the cloning of the V_{HH} in the Xho/Spe site of the Immuno PBS vector. However, the 3' end of the gene was not in phase with the identification "tag" and the stop codon of the vector. To achieve this, the construct was cut with Spe and the 4 base overhangs were filled in, using the Klenow fragment after which the vector was religated.

- The expression vector plasmid ipBS (immunopBS) (Stratagene) contains a pel B leader sequence which is used for immunoglobulin chain expression in E.coli under the promotor pLAC control, a ribosome binding site, and stop codons. In addition, it contains a sequence for a c-terminal decapeptide tag.

- E.coli JM101 harboring the ipBS-V_{HH}21 plasmid was grown in 1 l of TB medium with 100 µg/ml ampicillin and 0.1 % glucose at 32°C. Expression was induced by the addition of 1 mM IPTG (final concentration) at an OD₅₅₀ of 1.0. After overnight induction at 28°C, the

cells were harvested by centrifugation at 4.000 g for 10 min (4°C) and resuspended in 10 ml TES buffer (0.2 M Tris-HCL pH 8.0, 0.5 mM EDTA, 0.5 M sucrose). The suspension was kept on ice for 2 hours. Periplasmic proteins were removed by osmotic shock by addition of 20 ml TES buffer diluted 1:4 v/v with water, kept on ice for one hour and subsequently centrifugated at 12.000 g for 30 min. at 4°C. The supernatant periplasmic fraction was dialysed against Tris-HCl pH 8.8, NaCl 50mM, applied on a fast Q Sepharose flow (Pharmacia) column, washed with the above buffer prior and eluted with a linear gradient of 50 mM to 1 M NaCl in buffer.

Fractions containing the V_{HH} protein were further purified on a Superdex 75 column (Pharmacia) equilibrated with PBS buffer (0.01 M phosphate pH 7.2, 0.15 M NaCl). The yield of purified V_{HH} protein varies from 2 to 5 mg/l cell culture.

- Fractions were analyzed by SDS-PAGE(I). Positive identification of the camel V_{HH} antibody fragment was done by Western Blot analysis using antibody raised in rabbits against purified camel IgGH₃ and an anti-rabbit IgG-alkaline phosphatase conjugate (II).

As protein standards (Pharmacia) periplasmic proteins prepared from 1 ml of IPTG-induced JM101/ipBS V_{HH} 21 were used. Figure 8 shows: C,D:fractions from fast S Sepharose column chromatography (C:Eluted at 650 mM NaCl D:Eluted at 700 mM NaCl) E,F:fractions from Superdex 75 column chromatography.

As can be seen, the major impurity is eliminated by ionexchange chromatography and the bulk of the remaining impurities are eliminated by gel filtration.

Camel	γ_3 28Kd	-	L	P	G	G	P	S	V	F	V	F	P	P	P	K	P	K	D	V	L	S	I	X	G	X	P	-
Clone	# 72/1	-	L	P	G	G	P	S	V	F	V	F	P	T	K	P	K	D	V	L	S	I	S	G	R	P	-	-
Clone	# 72/4	-	L	P	G	G	P	S	V	F	V	F	P	P	K	P	K	D	V	L	S	I	S	G	R	P	-	-
Clone	# 72/29	-	L	L	G	G	P	S	V	F	I	F	P	P	K	P	K	D	V	L	S	I	S	G	R	P	-	-
Human	$\gamma_1\gamma_3$	-	L	L	G	G	P	S	V	F	L	F	P	P	K	P	K	D	T	L	M	I	S	R	T	P	-	-
CH2	γ_2	-	V	A	-	G	P	S	V	F	L	F	P	P	K	P	K	D	T	L	M	I	S	R	T	P	-	-
	γ_4	-	F	L	G	G	P	S	V	F	L	F	P	P	K	P	K	D	T	L	M	I	S	R	T	P	-	-

CH2 | CH3

		360	370																				
Camel	γ_3 14Kd	-	K	G	Q	T	R	E	P	Q	V	Y	T	L	A	P	X	R	L	E	L	-	-
Human	γ_1	-	K	G	Q	P	R	E	P	Q	V	Y	T	L	P	P	S	R	D	E	L	-	-
CH2/CH3	γ_2, γ_3	-	K	G	Q	P	R	E	P	Q	V	Y	T	L	P	P	S	R	E	E	M	-	-
	γ_4	-	K	G	Q	P	R	E	P	Q	V	Y	T	L	P	P	S	Q	E	E	M	-	-

Table 1

Comparison of the N terminal Camel CH2 and CH3 sequences with the translated cDNA sequences of Camel immunoglobulins and with the corresponding human γ sequences. (Numbering according to Kabat et al (1987)(7).

Primer Derived	10	20	30
	G G S V Q T G G S L R L S C E I S G L T F D		# 72/4
	G G S V Q T G G S L R L S C A V S G F S F S		# 72/3
	G G S E Q G G G S L R L S C A I S G Y T Y G		# 72/7
	G G S V Q P G G S L T L S C T V S G A T Y S		# 72/17
	G G S V Q A G G S L R L S C T G S G F P Y S		# 72/18
D V Q L V A S G G G S V G A G G S L R L S C T A S G D S F S			# 72/2
E V K L V E S G G G L V E P G G S L R L S C A T S G F T F S			Mouse $V_{HIII}A$
E V Q L L S G G G L V Q P G G S L R L S C A A S G F T F S			Human V_{HIII}

Table 2:

A comparison of N Terminal Fr I regions of Camel V_{HH} with a Human V_{HIII} subgroup protein and a mouse $V_{HIII}A$ subgroup protein.

The invariable subgroup specific residues are grayed.

	FrameWork 4	J Genes
Human	W G Q G T L V T V S S	J1, J4, J5
	W G R G T L V T V S S	J2
	W G Q G T T V T V S S	J6
	W G Q G T M V T V S S	J3
Murine	W G Q G T T L T V S S	J1
	W G Q G T L V T V S S	J2
	W G Q G T S V T V S A	J3
	W G A G T T V T V S S	J4
		cDNA Clones
Camel	W G Q G T Q V T V S S	Clones
	W G Q G T Q V T V S S	# 72/19 = # 72/3
	W G Q G T L V T V S S	1 Clone
	W G R G T Q V T V S S	# 72/24
	W G Q G T H V T V S S	# 72/21
	W G Q G I Q V T A S S	# 72/16

Table 3

Comparison of some Framework 4 residues found in the Camel V_HH region with the Framework 4 residues corresponding to the consensus region of the Human and Mouse J minigenes.

	MOUSE VH3 sequence	
D Y Y G S S 95	100 a b c d e f g h i j k	110
1	A L Q L M H L G P G I S Q G Y G X	V S S
2	V P A S G S G T A G C E L P L G G	V S S L
3	F C Y S G S G T A G C E L P L G G	V S S
4	E L S G S G T A G C E L P L G G	V S S
7	D W K Y M G T R V S I Y M P Y C N L D P N Y A T L A P W S V	V S S
9	R L K D F S S I Y M P Y C N L D P N Y A T L A P W S V	V S S
11	Q K R F S S I Y M P Y C N L D P N Y A T L A P W S V	V S S
13	G S R F S S I Y M P Y C N L D P N Y A T L A P W S V	V S S
16	A D P S I Y M P Y C N L D P N Y A T L A P W S V	V S S
17	G S R F S S I Y M P Y C N L D P N Y A T L A P W S V	V S S
18	D S S F Y W Y G C N L D P N Y A T L A P W S V	V S S
19	T E I L A G G W Y C N L D P N Y A T L A P W S V	V S S
20	N Q L T E M G A C D A R G I G L	V S S
21	R L T E M G A C D A R G I G L	V S S
24	D G W T P C H L L	V S S
25	D S Y P I A D M C S	V S S
27	V E Y P I A D M C S	V S S
29	V E Y P I A D M C S	V S S

Random Sample

18 different
camel VHH region

CDR3

Human and mouse	size range 0-19 aa	over 600 entries
Camel	8-24 aa	18 entries

TABLE 4

55/1

human C _H 1		hinge	C _H 2
human gamma 3	KVDKRV	ELKTPLGDTTHTCPRCP	
		<u>EPKCSDTTPPPCPRCP</u>	
		<u>EPKSCDTTPPPCPRCP</u>	APPELLGG PSVFLFP
human gamma 1	KVDKK	— AEPKSCDKTHTCPPCP	APPELLGG PSVFLFP
human gamma 2	KVKVTV	— ERKCCVVECPPCP	APPVAG - PSVFLFP
human gamma 4	KVDKRV	— ESKYGPCCPSCP	APPELLGG PSVFLFP

TABLE 5(2)

10	20	40
EVQLVESGGG	LVQPGGSLRL SCAASG	CDR1: NVRQA PGKGLEWVS CDR2:
GG SVQGGGSLRL	SCAISG	CDR1: WFRREG PGKEREGLA CDR2:
GG SVQAGGSLRL	SCASSS	CDR1: WYRQA PGKEREFEVS CDR2:

70	80	90	110
RFTIS RDNSKNTLYL	QMNSLRAEDTAVY YCAR	CDR3: WGQGTQVT	VSS
RFTIS QDSTLKTMYL	LMNNLKPEDTGT YCAA	CDR3: WGQGTQVT	VSS
RFTIS QDSAKNTVYL	QMNSLKPEDTAMY YCKI	CDR3: WGQGTQVT	VSS

camel	camel V _H	hinge	C _H 2
	WGQGTQVT VSS	GTNEVCKCPKCP	APELPGG PSVFVFP
	WGQGTQVT VSS	EPKIPQPKPQPKPQ	QPKPKPQ
		KPEPECTCPKCP	APELLGG PSVFIFP

TABLE 5(1)

SUBSTITUTE SHEET

REFERENCES

1. Ward, E.S., Güssow, D., Griffiths, A.D., Jones, P.T. and Winter G. Nature 341, 544-546 (1989).
2. Ungar-Waron H., Eliase E., Gluckman A. and Trainin Z. Isr. J. Vet. Med., 43, 198-203 (1987).
3. Bajyana Songa E. and Hamers R., Ann. Soc. Belge Méd. trop., 68, 233-240 (1988).
4. Edelman G.M., Olins D.E., Gally J.A. and Zinder N.D., Proc. Nat. Acad. Sc., 50, 753 (1963).
5. Franek F. and Nezlin R.S., Biokhimiya, 28, 193 (1963).
6. Roitt I.M., Brostoff J. and Male D.K., Immunology, Gower Med. Pub. London. New-York, p.9.2. (1985).
7. Schiffer M., Girling R.L., Ely K.R. and Edmundson B., Biochemistry, 12, 4620-4631 (1973).
8. Fleischman J.B., Pain R.H. and Porter R.R., Arch. Biochem. Biophys, Suppl. 1, 174 (1962).
9. Roholt O., Onoue K. and Pressman D., PNAS 51, 173-178 (1964).
10. Seligmann M., Mihaesco E., Preud'homme J.L., Danon F. and Brouet J.C., Immunological Rev., 48, 145-167 (1979).
11. Hendershot L., Bole D., Köhler G. and Kearney J.F., The Journal of Cell Biology, 104, 761-767 (1987).
12. Hendershot L.M., The Journal of Cell Biology, 111, 829-837 (1990).

13. Hamers-Casterman, C., E. Wittouck, W. Van der Loo and R. Hamers, Journal of Immunogenetics, 6, 373-381 (1979).
14. Applied Biosystems - Ethanol Precipitation of Electro Eluted Electrodialysed Sample. Issue n° 27.
15. Maniatis, T. E.F. Fritsch and J. Sambrook, Molecular Cloning. A Laboratory Manual (1988).
16. Sastry et al., PNAS, 86, 5728, (1989).
17. Sanger, F., S. Nicklen and A.R. Coulson, Proc. Natl. Acad. Sci., U.S.A., 74, 5463-5467 (1977).
18. Kabat E.A., Tai Te Wu, M. Reid-Miller, H.M. Perry and K.S. Gottesman, U.S. Dpt of Health and Human Services, Public Health Service, National Institutes of Health (1987).
19. Valentine, R.C. and N.M. Geen, J.M.B., 27, 615-617 (1967).

C L A I M S

1. Immunoglobulin characterized in that it comprises two heavy polypeptide chains sufficient for the formation of a complete antigen binding site or several antigen binding sites, this immunoglobulin being further devoid of light polypeptide chains.
2. Immunoglobulin according to claim 1, characterized in that it comprises two heavy polypeptide chains sufficient for the formation of a complete antigen binding site or several antigen binding sites, this immunoglobulin being further devoid of light polypeptide chains and further characterized by the fact that it is the product of the expression in a prokaryotic or in a eukaryotic host cell, of a DNA or of a cDNA having the sequence of an immunoglobulin devoid of light chains as obtainable from lymphocytes or other cells of Camelids.
3. Immunoglobulin according to claim 1 or claim 2, characterized in that the amino acid sequence of its variable region contains in position 45 an amino acid which is different from a leucine, or proline or glutamine residue.
4. Immunoglobulin according to anyone of claims 1 to 3, characterized in that its heavy polypeptide chains are devoid of a so-called first domain in their constant region (C_H1).
5. Immunoglobulin according to anyone of claims 1 to 4, characterized in that it comprises an antigen binding site or several antigen binding sites and especially in that each variable region of each heavy chain contains at least one antigen binding site.
6. Immunoglobulin according to anyone of claims 1 to 5, characterized in that it is a type G immunoglobulin of class 2 (IgG2) or it is a type G immunoglobulin of class 3 (IgG3).

7. Immunoglobulin according to anyone of claims 1 to 6, characterized in that it is a Camelid immunoglobulin.

8. Immunoglobulin according to anyone of claims 1 to 7 as obtainable by purification from the serum of Camelids, characterized in that :

- it is not adsorbed by chromatography on Protein G Sepharose column,
- it is adsorbed by chromatography on Protein A Sepharose column,
- it has a molecular weight of around 100 Kd after elution with a pH 4.5 buffer (0.15 M NaCl, 0.58% acetic acid adjusted to pH 4.5 by NaOH),
- it consists of heavy $\gamma 2$ polypeptide chains of a molecular weight of around 45 Kd preferably 46 Kd after reduction.

9. Immunoglobulin according to anyone of claims 1 to 7, as obtainable by purification from the serum of Camelids is characterized in that the immunoglobulin :

- is adsorbed by chromatography on a Protein A Sepharose column,
- has a molecular weight of around 100 kd after elution with a pH 3.5 buffer (0.15 M NaCl, 0.58% acetic acid),
- is adsorbed by chromatography on a Protein G Sepharose column and eluted with pH 3.5 buffer (0.15 M NaCl, 0.58% acetic acid).
- consists of heavy $\gamma 3$ polypeptide chains of a molecular weight of around 45 Kd in particular between 43 and 47 kd after reduction.

10. Immunoglobulin according to anyone of claims 1 to 9, characterized in that

- it comprises 4 frameworks in its variable region, which frameworks comprise an amino-acid sequence selected from the following sequences :

for the framework 1 domain

60

G G S V Q T G G S L R L S C E I S G L T F D
 G G S V Q T G G S L R L S C A V S G F S F S
 G G S E Q G G G S L R L S C A I S G Y T Y G
 G G S V Q P G G S L T L S C T V S G A T Y S
 G G S V Q A G G S L R L S C T G S G F P Y S
 G G S V Q A G G S L R L S C V A G F G T S
 G G S V Q A G G S L R L S C V S F S P S S

for the framework 4 domain

W G Q G T Q V T V S S
 W G Q G T L V T V S S
 W G Q G A Q V T V S S
 W G Q G T Q V T A S S
 R G Q G T Q V T V S L

and/or,

for the CDR3 domain

A L Q P G G Y C G Y G X - - - - - C L
 V S L M D R I S Q H - - - - - G C
 V P A H L G P G A I L D L K K Y - - - - - K Y
 F C Y S T A G D G G S G E - - - - - M Y
 E L S G G S C E L P L L F - - - - - D Y
 D W K Y W T C G A Q T G G Y F - - - - - G Q
 R L T E M G A C D A R W A T L A T R T F A Y N Y
 Q K K D R T R W A E P R E W - - - - - N N
 G S R F S S P V G S T S R L E S - S D Y - - N Y
 A D P S I Y Y S I L X I E Y - - - - - K Y
 D S P C Y M P T M P A P P I R D S F G W - - D D
 T S S F Y W Y C T T A P Y - - - - - N V
 T E I E W Y G C N L R T T F - - - - - T R
 N Q L A G G W Y L D P N Y W L S V G A Y - - A I
 R L T E M G A C D A R W A T L A T R T F A Y N Y
 D G W T R K E G G I G L P W S V Q C E D G Y N Y
 D S Y P C H L L - - - - - D V
 V E Y P I A D M C S - - - - - R Y

and/or,

- in that its constant region comprises C_H2 and C_H3 domains comprising an amino-acid sequence selected from the following sequences :

for the C_H2 domain:

APELLGGPTVFIFPPKPKDVLSITLTP
 APELPGGPSVFVFPTKPKDVLSISGRP
 APELPGGPSVFVFPPKPKDVLSISGRP
 APELLGGPSVFIFPPKPKDVLSISGRP

for the C_H3 domain:

GQTREPQVYTLA
 GQTREPQVYTLAPXRLEL
 GQPREPQVYTLPPSRDEL
 GQPREPQVYTLPPSREEM
 GQPREPQVYTLPPSQEEM

and/or,

- in that its hinge region comprises from 0 to 50 amino-acids, especially in that its hinge region comprises an amino-acid sequence selected from the following sequences :

GTNEVCKCPKCP

or,

EPKIPQPKPQPKPQPKPQPKPQPKPEPECTCPKCP

11. Immunoglobulin according to anyone of claims 1 to 10, characterized in that it is encoded by a sequence selected among those represented on figure 7.

12. Fragment of an immunoglobulin according to anyone of claims 1 to 11, characterized in that it is selected from the following group :

- a fragment corresponding to one heavy polypeptide chain of an immunoglobulin devoid of light chains,
- fragments obtained by enzymatic digestion of the immunoglobulins of the invention, especially those obtained by partial digestion with papain leading to the Fc fragment (constant fragment) and leading to FV_{HH}h fragment (containing the

antigen binding sites of the heavy chains) or its dimer $F(V_{HH}h)_2$, or a fragment obtained by further digestion with papain of the Fc fragment, leading to the Fc' fragment corresponding to the C-terminal part of the Fc fragment,

- homologous fragments obtained with other proteolytic enzymes,
- a fragment of at least 10 preferably 20 amino acids of the variable region of the immunoglobulin, or the complete variable region, especially a fragment corresponding to the isolated V_{HH} domains or to the V_{HH} dimers linked to the hinge disulphide,
- a fragment corresponding to the hinge region of the immunoglobulin, or to at least 6 amino acids of this hinge region,
- a fragment of the hinge region comprising a repeated sequence of Pro-X,
- a fragment corresponding to at least 10 preferably 20 amino acids of the constant region or to the complete constant region of the immunoglobulin.

13. Immunoglobulin according to anyone of claims 1 to 12, characterized in that all or a part of its constant region is replaced by all or part of the constant region of a human antibody.

14. Immunoglobulin according to any one of claims 1 to 13, obtainable in prokaryotic cells, especially in E.coli cells by a process comprising the steps of :

- a) cloning in a Bluecript vector of a DNA or cDNA sequence coding for the V_H domain of an immunoglobulin devoid of light chain obtainable for instance from lymphocytes of Camelids,
- b) recovering the cloned fragment after amplification using a 5' primer containing an Xho site and a 3' primer containing the Spe site having the following sequence

TC TTA ACT AGT GAG GAG ACG GTG ACC TG,

- c) cloning the recovered fragment in phase in the immuno PBS vector after digestion of the vector with Xho and Spe restriction enzymes,
- d) transforming host cells, especially E.coli by transfection with the recombinant immuno PBS vector of step c,
- e) recovering the expression product of the V_{HH} coding sequence, for instance by using antibodies raised against the dromadary V_{HH} domain.

15. Hetero-specific immunoglobulins according to any one of claims 1 to 13 obtainable by a process comprising the steps of:

- obtaining a first DNA or cDNA sequence coding for a V_{HH} domain or part thereof having a determined specificity against a given antigen and comprised between Xho and Spe sites,
- obtaining a second DNA or cDNA sequence coding for a V_{HH} domain or part thereof, having a determined specificity different from the specificity of the first DNA or cDNA sequence and comprised between the Spe and EcoRI sites,
- digesting an immuno PBS vector with EcoRI and XhoI restriction enzymes,
- ligating the obtained DNA or cDNA sequences coding for V_{HH} domains, so that the DNA or cDNA sequences are serially cloned in the vector,
- transforming a host cell, especially E.coli cell by transfection, and recovering the obtained immunoglobulins.

16. Immunoglobulin according to any one of claims 1 to 13, obtainable by a process comprising the steps of:

- obtaining a DNA or cDNA sequence coding for a V_{HH} domain or part thereof, having a determined specific antigen binding site,

- amplifying the obtained DNA or cDNA, using a 5' primer containing an initiation codon and a HindIII site, and a 3' primer containing a termination codon having a XhoI site,
 - recombining the amplified DNA or cDNA into the HindIII (position 2650) and XhoI (position 4067) sites of a plasmid pMM984,
 - transfecting permissive cells especially NB-E cells with the recombinant plasmid,
 - controlling the expression, for instance by an ELISA assay with antibodies directed against a region of a V_{HH} domain and recovering the obtained products.
17. Immunoglobulins according to claim 16, obtainable by a process comprising the further cloning of a second DNA or cDNA sequence having another determined antigen binding site, in the pMM984 plasmid.
18. Immunoglobulin according to any one of claims 11 to 17, characterized in that it is obtainable by a process wherein the vector is Yep 52 and the transformed recombinant cell is a yeast especially S.cerevisiae.
19. Immunoglobulin according to any one of claims 11 to 17, characterized in that it is obtainable by a process wherein the vector is a vector appropriate for expression in plant cells, for example pMon530, and the transformed recombinant cells are plant cells.
20. Immunoglobulin according to any one of claims 16 to 17, characterized in that it has a catalytic activity, especially in that it is directed against an antigen mimicking an activated state of a given substrate, these immunoglobulins having for instance been modified at the level of their catalytic site by random or directed mutagenesis.
21. Nucleotide sequence, characterized in that it codes for all or part of an immunoglobulin according to anyone of claims 1 to 20, which immunoglobulins

comprise a peptide sequence selected from the following:

VTVSSGTNEVCKCPKCPAPELPGGPSVVFVFP,
 VTVSSEPKIPQPKPQPKPQPKPQPKPEPECTCPKCPAPELLGGPSVFIFP
 GTNEVCKCPKCP
 APELPGGPSVVFVFP
 EPKIPQPKPQPKPQPKPQPKPQPKPEPECTCPKCP
 APELLGGPSVFIFP
 APELLGGPTVFIFPPKPKDVLSITLTP
 APELPGGPSVVFVFTKPKDVLSISGRP
 APELPGGPSVVFVFPKPKDVLSISGRP
 APELLGGPSVFIFPPKPKDVLSISGRP
 GQTREPQVYTLA
 GQTREPQVYTLAPXRLEL
 GQPREPQVYTLPPSRDEL
 GQPREPQVYTLPPSREEM
 GQPREPQVYTLPPSQEEM
 G G S V Q T G G S L R L S C E I S G L T F D
 G G S V Q T G G S L R L S C A V S G F S F S
 G G S E Q G G G S L R L S C A I S G Y T Y G
 G G S V Q P G G S L T L S C T V S G A T Y S
 G G S V Q A G G S L R L S C T G S G F P Y S
 G G S V Q A G G S L R L S C V A G F G T S
 G G S V Q A G G S L R L S C V S F S P S S
 W G Q G T Q V T V S S
 W G Q G T L V T V S S
 W G Q G A Q V T V S S
 W G Q G T Q V T A S S
 R G Q G T Q V T V S L
 and/or,

A L Q P G G Y C G Y G X - - - - - C L
 V S L M D R I S Q H - - - - - G C
 V P A H L G P G A I L D L K K Y - - - - - K Y
 F C Y S T A G D G G S G E - - - - - M Y
 E L S G G S C E L P L L F - - - - - D Y
 D W K Y W T C G A Q T G G Y F - - - - - G Q

```

R L T E M G A C D A R W A T L A T R T F A Y N Y
Q K K D R T R W A E P R E W - - - - - N N
G S R F S S P V G S T S R L E S - S D Y - - N Y
A D P S I Y Y S I L X I E Y - - - - - K Y
D S P C Y M P T M P A P P I R D S F G W - - D D
T S S F Y W Y C T T A P Y - - - - - N V
T E I E W Y G C N L R T T F - - - - - T R
N Q L A G G W Y L D P N Y W L S V G A Y - - A I
R L T E M G A C D A R W A T L A T R T F A Y N Y
D G W T R K E G G I G L P W S V Q C E D G Y N Y
D S Y P C H L L - - - - - D V
V E Y P I A D M C S - - - - - R Y

```

22. Nucleotide sequence characterized in that it codes for an immunoglobulin according to anyone of claims 1 to 20, in that it comprises a sequence selected from those represented on figure 7.

23. Process for the preparation of a monoclonal antibody according to anyone of claims 1 to 20, directed against a determined antigen, the antigen binding site of the antibody consisting of heavy polypeptide chains and which antibody is further devoid of light polypeptide chains, which process comprises :

- immortalizing lymphocytes, obtained for example from the peripheral blood of Camelids previously immunized with a determined antigen, with an immortal cell and preferably with myeloma cells, in order to form a hybridoma,
- culturing the immortalized cells formed and recovering the cells producing the antibodies having the desired specificity.

24. Process for the preparation of antibodies directed against determined antigens, comprising the steps of :

- cloning into vectors, especially into phages and more particularly filamentous bacteriophages, DNA

or cDNA sequence obtained from lymphocytes of Camelids previously immunized with determined antigens, capable of producing an immunoglobulin according to anyone of claims 1 to 20,

- transforming prokariotic cells with the above vectors in conditions allowing the production of the antibodies,
- selecting the appropriate antibody by subjecting the transforming cells to antigen-affinity selection,
- recovering the antibodies having the desired specificity.

25. Process according to claim 24, wherein the cloning vector is a plasmid or a eukaryotic virus and the transformed cell is a eukaryotic cell, especially a yeast cell, mammalian cell, plant cell or protozoair cell.

26. Process according to claim 24, wherein the cloning vector is a plasmid capable of expressing the immunoglobulin in the bacterial membrane.

27. Process according to claim 24, wherein the cloning vector is a plasmid capable of expressing the immunoglobulin as a secreted protein.

28. Immunoglobulin according to anyone of claims 1 to 20, characterized in that it is directed against an antigen such as one of a bacteria, a virus, a parasite, or against a protein, hapten, carbohydrate or nucleic acid.

29. Immunoglobulin according to anyone of claims 1 to 20 characterized in that it is directed against an immunoglobulin idiotype.

30. Immunoglobulin according to anyone of claims 1 to 20 characterized in that it is directed against a cellular receptor or membrane protein.

31. Immunoglobulin according to anyone of claims 1 to 20, characterized in that it has a catalytic activity.

32. Immunoglobulin according to anyone of claims 1 to 20, or a fragment according to claim 12, characterized in that it is conjugated with a toxin.

33. Use for a fragment comprising a repeated sequence Pro-X, X being any amino-acid and preferably Gln, Lys or Glu, the sequence containing advantageously at least 3 repeats of Pro-X and especially a fragment composed of a 12-fold repeat of the sequence Pro-X, for coupling protein domains or a protein and a ligand.

34. Use of the hinge region or of a fragment of the hinge region of an immunoglobulin according to anyone of claims 1 to 20, for coupling protein domains or a protein and a ligand.

35. Immunoglobulin according to anyone of claims 1 to 20, characterized in that it is a heterospecific antibody.

36. Recombinant vector characterized in that it comprises a nucleotide sequence according to claim 21 or claim 22, and in that it is a plasmid, a phage especially a bacteriophage, a virus, a YAC, a cosmid.

37. Recombinant cell or organism characterized in that it is modified by a vector according to claim 36.

38. A cDNA library composed of nucleotide sequences coding for a heavy-chain immunoglobulin according to anyone of claims 1 to 20, such as obtained by performing the following steps:

a) treating a sample containing lymphoid cells, especially peripheral lymphocytes, spleen cells, lymph nodes or another lymphoid tissue from a healthy animal, especially selected among the Camelids, in order to separated the B-lymphocytes,

b) separating polyadenylated RNA from the other nucleic acids and components of the cells,

c) reacting the obtained RNA with a reverse transcriptase in order to obtain the corresponding cDNA,

d) contacting the obtained cDNA with 5' primers corresponding to mouse V_H domain of four-chain immunoglobulins, which primer contains a determined restriction site, for example an XhoI site and with 3' primers corresponding to the N-terminal part of a C_H2 domain,

e) amplifying the DNA,

f) cloning the amplified sequence in a vector, especially in a bluescript vector,

g) recovering the clones hybridizing with a probe corresponding to the sequence coding for a constant domain from an isolated heavy-chain immunoglobulin.

39. A modified 4-chain immunoglobulin or a fragment thereof, the V_H regions of which has been partially replaced by specific sequences or amino acids of heavy chain immunoglobulins according to anyone of claims 1 to 20.

40. A modified 4-chain immunoglobulin or a fragment thereof, according to claim 39, wherein the leucine, proline or glutamine in position 45 of the V_H regions has been replaced by other amino acids and preferably by arginine, glutamic acid or cysteine.

41. A modified 4-chain immunoglobulin or a fragment thereof, in which the CDR loops of the region are linked to other parts of the V region by the introduction of paired cysteines, in particular in which the CDR_3 loop is linked to the FW_2 or CDR_1 and more especially where the cysteine of the CDR_3 of the V_H is linked to a cysteine in position 31 or 33 of FW_2 or in position 45 of CDR_2 .




FIGURE 1A

SUBSTITUTE SHEET




FIGURE 1B




FIGURE 1C




FIGURE 2A






FIGURE 2B




Prot. A	Ig1	Ig2	Ig3	Tot.Ser	Ig1	Ig2	Ig3	Tot.Ser	
Control	T. evansi Infected				Healthy				
Counts/5ul	65	1258	1214	2700	2978	147	157	160	107

FIGURE 3A



Ig1 Ig2 Ig3 Ig1 Ig2 Ig3
Healthy T. evansi infected

FIGURE 3B



Ig1 Ig2 Ig3
T. evansi infected
Ponceau Red

FIGURE 3C




FIGURE 4A




FIGURE 4B

Analysis of IgG₃ Papain Fragments by SDS - PAGE

SUBSTITUTE SHEET

Model for Camel IgG₃




Fig 5: Schematic representation of Camel IgG₃ model.




Fig 6: A schematic representation of Camel immunoglobulins IgG₁, the putative IgG₂ and IgG₃. The large (Pro-X)₁₂ of the putative IgG₂ molecule can be modelled into a 6 aa repeat

10 / 14

```

DR01006 C-----TCGAG---TCTGGGGGAGG
DR27006 C-----TCGAG---TCTGGGGGAGG
DR03006 C-----AGGTGA-----AACTGCTCGAG---TCTGGAGGAGG
DR11006 C-----TCGAG---TCTGGGGGAGG
DR24006 C-----AGGTGA-----AACTGCTCGAG---TCTGGGGGAGG
DR16006 C-----TCGAG---TCTGGAGGAGG
DR19006 C-----TCGAG---TCTGGAGGAGG
DR07006 C-----TCGAG---TCTGGGGGAGG
DR16006 C-----TCGAG---TCTGGGGGAGG
DR20006 C-----TCGAG---TCAGGGGGAGG
DR25006 C-----TCGAG---TCTGGGGGAGG
DR20006 C-----TCGAG---TCTGGAGGAGG
DR21006 C-----TCGAG---TCTGGGGGAGG
DR09006 C-----AGGTGA-----AACTGCTCGAG---TCTGGGGGAGG
DR17006 C-----TCGAG---TCTGGGGGAGG
DR13006 C-----TCGAG---TCAGGGGGAGG
DR02006 CTCGAGTCAGGTGTCCGGTCTGATGTGCAGCTGGTGGCGTCTGGGGGAGG

```

```

DR01006 ATCGGTGCAGGCTGGAGGGTCTCTGAGACTCTC--GTGCG-CAGCCTCTG
DR27006 CTCGGTGCAGGCTGGAGGGTCTCTGAGACTCTCCTGTGCATCTTCTTCTA
DR03006 CTCGGTGCAGACTGGAGGATCTCTGAGACTCTCCTGTGCAGT--C-TCTG
DR11006 GTCGGTGCAGGCTGGAGGGTCTCTGAGACTCTCCTGTAATGT--C-TCTG
DR24006 GTCGGTGCAGGCTGGAGGGTCTCTGAGACTCTCCTGTAATGT--C-TCTG
DR16006 CTCGGCGCAGGCTGGAGGATCTCTGAGACTCTCCTGTGCAGC--CCACGG
DR19006 CTCGGTTCAGGCTGGAGGGTCCCTTAGACTCTCCTGTGCAGC--C-TCTG
DR07006 CTCGGTGCAGGGTGGAGGGTCTCTGAGACTCTCCTGTGCAA---TCTCTG
DR16006 CTCGGTGCAGGCTGGAGGGTCTCTGAGACTCTCCTGTACAG---GCTCTG
DR20006 CTCGGTACAGGTTGGAGGGTCTCTGAGACTCTCCTGTGTAG---CCTCTA
DR25006 CTCGGTACAACTGGAGGGTCTCTGAGACTCTCTTGCG---AAATCTCTG
DR20006 CTCGGTGCAGGCTGGAGGGTCTCTGAGACTCTCCTGTG---TAGCCTCTG
DR21006 CTCGGTGCAGGTTGGAGGGTCTCTGAAACTCTCCTGTAAAAT---CTCTG
DR09006 CTCGGTGCAGGCTGGGGGGTCTCTGACACTCTCTTGTG---TATACAC--
DR17006 CTCGGTCCAACCTGGAGGATCTCTGACACTCTCCTGTACAGTT---TCTG
DR13006 CTCGGTGGAGGCTGGAGGGTCTCTGAGACTCTCCTGTACAG---CCTCTG
DR02006 CTCGGTGCAGGCTGGAGGGTCTCTGAGACTCTCCTGTACAG---CCTCTG

```

```

DR01006 GA--TACAGTAATT---GTCCCCTCACTTG-GAGCTGGTATCGCCAGTTT
DR27006 AA--TATATGCCTT---GCACCTACGACAT-GACCTGGTACCGCCAGGCT
DR03006 GA--TTCTCCTTTA---GTACCAGTTGTAT-GGCCTGGTTCCGCCAGGCT
DR11006 GC--TCTCCAGTA---GTACTTATTGCCT-GGGCTGGTTCCGCCAGGCT
DR24006 GC--TCTCCAGTA---GTACTTATTGCCT-GGGCTGGTTCCGCCAGGCT
DR16006 GA--TTCCGC-TCA---ATGGTTACTACAT-CGCCTGGTTCCGTCAGGCT
DR19006 AC--TACACATCA---CTGATTATTGCAT-GGCCTGGTTCCGCCAGGCT
DR07006 GA--TACACGTACG---GTAGCTTCTGTAT-GGGCTGGTTCCGCCAGGCT
DR16006 GA--TTCCCCTATA---GTACCTTCTGTCT-GGGGTGGTTCCGCCAGGCT
DR20006 CT--CACACCGACA---GTAGCACCTGTAT-AGGCTGGTTCCGCCAGGCT
DR25006 GA--TTGACTTTTG---ATGATTCTGACGT-GGGGTGGTACCGCCAGGCT
DR20006 GA--TTCAATTTTCG---AAACTTCTCGTAT-GGCGTGGTACCGCCAGACT
DR21006 GAGGTACCCAGATCGTGTTCCTAAATCTTTGGCCTGGTTCCGCCAGGCT

```

FIGURE 7(1)

SUBSTITUTE SHEET

DR09006 -----CAACGATACTGGGACCA-----TGGGATGGTTTCGCCAGGCT
 DR17006 --GGGCCACCTACA---GTGACTACAGTATTGGA-TGGATCCGCCAGGCT
 DR13006 G-----ATACGTAT-CCT----CTATGGCCTGGTTCCGCCAGGTT
 DR02006 GAGA---CAGTTTCAGTAGATT--TGCCATGTCTTGGTTCCGCCAGGCT

DR01006 CCAGGAACGGAGCGCGAGTTTCGTCTCCAGTATGGATCCGGATGGAAATAC
 DR27006 CCAGGCAAGGAGCGCGAATTTGTCTCAAGTATAAATATTGATGGTAAGAC
 DR03006 TCAGGAAAGCAGCGTGAGGGGGTCGCAGCCATTAATAGTGGCGGTGGTAG
 DR11006 CCAGGGAGGGAGCGTGAGGGGGTCACAGCGATTAA-----CACTGATGG
 DR24006 CCAGGGAAGGAGCGTGAGGGGGTCACAGCGATTAA-----CACTGATGG
 DR16006 CCTGGGAAGGGCGTGAGGGGGTCGCAACAATTAATGGTGGTCG-----
 DR19006 CCAGGGAAGGAGCGTGAATTGGTTCGCAGCGATTCAAGTTGTCCGTAGTGA
 DR07006 CCAGGCAAGGAACGTGAGGGGATCGCAACTATTCTTAATGGTGGTACTAA
 DR16006 CCAGGGAAGGAGCGTGAGGGGGTCGCGGGTATTAATAGTGCAGGAGGTAA
 DR20006 CCAGGGAAGGAGCGCGAGGGGGTCGCAAGTATATATTTTGGTGATGGTGG
 DR25006 CCAGGGGATCAGTGCAAATTTGGTCTCAGGTATTCTGAGTGATGGTACT-C
 DR20006 CCAGGAAATGTGTGTGAGTTGGTCTCAAGTATTTACAGTGATGG-----
 DR21006 CCAGAGAAGGAGCGCGAGGGGATCGCAGTTCCTTCGACTAAGGATGGTAA
 DR09006 CCAGGGAAGAGTGCGAAAGGGTCGCGCATATTACGCTGATGGTATGA-
 DR17006 CCAGGGAAGGACCGTGAAGTAGTCGCAGCCGCTAATACTGGTG-----
 DR13006 CCAGGCGAGGAGCGCGAGGGGGTCGCGTTTGTTCAAACGG-----
 DR02006 CCAGGGAAGGAGTGCGAATTGGTCTCAAGCATTCAAAGTAATGGAAGGAC

DR01006 CAAGTACA-----CATACTCCGTGAAGGGCCGCTTCACC
 DR27006 AACATACG-----CAGACTCCGTGAAGGGCCGATTTCACC
 DR03006 GACATACTA-CAACACATATGTCGCCGAGTCCGTGAAGGGCCGATTTCGCC
 DR11006 CAGTATCAT-ATACGCA-----GCCGACTCCGTGAAGGGCCGATTTCACC
 DR24006 CAGTGTCAT-ATACGCA-----GCCGACTCCGTGAAGGGCCGATTTCACC
 DR16006 -----CGA-CGTCACATACTACGCCGACTCCGTGACGGGGCCGATTTACC
 DR19006 TACT--CGC-C-TCACAGACTACGCCGACTCCGTGAAGGGACGATTTCACC
 DR07006 -----CACATACTATGCCGACTCCGTGAAGGGCCGATTTCACC
 DR16006 -----TACTTACTATGCCGACGCCGTGAAGGGCCGATTTCACC
 DR20006 -----TACGAATTATCGCGACTCCGTGAAGGGCCGATTTCACC
 DR25006 CATATACAAAGAGTGGAGACTATGCTGAGTCTGTGAGGGGCCGGTTACC
 DR20006 CA-AAACATACTACGTCGACC--GCA-----TGAAGGGCCGATTTCACC
 DR21006 GA-----CATTCATGCGGACTCCGTGAAGGGCCGATTTCACC
 DR09006 -----CCTTCATTGATGAACCCGTGAAGGGCCGATTTCAGG
 DR17006 -----CGACTAGTAAATCTACGTCGACTTTGTGAAGGGCCGATTTCACC
 DR13006 --CTGACAAT-AGTGCAATTATATGGCGACTCCGTGAAGGGCCGATTTCACC
 DR02006 AACTGA-----GGCCGATTCCGTGCAAGGGCCGATTTCACC

DR01006 ATGTCCCGAGGCAGCACCGAGTACACAGTATTTCTGCAAATGGACAATCT
 DR27006 ATCTCCCAAGACAGCGCCAAGAACACGGTGATCTGCAGATGAACAGCCT
 DR03006 ATCTCCCAAGACAACGCCAAGACCACGGTATATCTTGATATGAACAACCT
 DR11006 ATCTCCCAAGACACCGCCAAGGAAACGGTACATCTCCAGATGAACAACCT
 DR24006 ATCTCCCAAGACACCGCCAAGAAAACGGTATATCTCCAGATGAACAACCT
 DR16006 ATCTCCCGAGACAGCCCCAAGAATACGGTGATCTGCAGATGAACAGCCT
 DR19006 ATCTCCCAAGGCAACACCAAGAACACAGTGAATCTGCAAATGAACAGCCT
 DR07006 ATCTCCCAAGACAGCACGTTGAAGACGATGTATCTGCTAATGAACAACCT
 DR16006 ATCTCCCAAGGGAATGCCAAGAATACGGTGTCTCTGCAAATGGATAACTT
 DR20006 ATCTCCCAACTCAACGCCCCAGAACACAGTGATCTGCAAATGAACAGCCT
 DR25006 ATCTCCAGAGACAACGCCAAGAATGATATACCTTCAAATGAACAGCCT
 DR20006 ATTTCTAGAGAGAATGCCAAGAATACATTGTATCTACAACCTGAGCGGCCT
 DR21006 ATCTTCTTAGATAATGACAAGACCACTTTCTCCTTACAACCTTGATCGACT
 DR09006 ATCTCCCGAGACAACGCCCCAGAAAACGTTGTCTTTGCGAATGAATAGTCT

FIGURE 7(2)

SUBSTITUTE SHEET

12 / 14

DR17006 ATTTCCCAAGACAACGCCAAGAATACGGTATATCTGCAAATGAGCTTCCT
 DR13006 ATCTCCCACGACAACGCCAAGAACACGCTGTATCTGCAAATGCGCAACCT
 DR02006 ATCTCCCAGACAATTCCAGGAACACAGTGTATCTGCAAATGAACAGCCT

DR01006 GAAACCTGAGGACACGGCGATGTATTACTGTAAAAC-A---GCCCTAC--
 DR27006 GAAACCTGAGGACACGGCGATGTATTACTGTAAAAT-A---GA--TTC--
 DR03006 AACCCCTGAAGACACGGCTACGTATTACTGTGCGGCGG---TCCCAGCCC
 DR11006 GCAACCTGAGGATACGGCCACCTATTACTGCGCGGCAA---GACTGACGG
 DR24006 GCAACCTGAGGATACGGCCACCTATTACTGCGCGGCAA---GACTGACGG
 DR16006 GAAACCTGAGGACACGGCCATCTACTTCTGTGCAGCAG---G-----CTC
 DR19006 GACACCTGAGGACACGGCCATCTACAGTTGTGCGGCAA---C-----CAG
 DR07006 GAAACCTGAAGACACGGGCACCTATTACTGTGCTG-CA---GAACTAAGT
 DR16006 GAAACCTGAGGACACGGCCATCTATTACTGCGCGG-CG---GATAGTCCA
 DR20006 GAAACCTGAGGACAGCGCCATGTACTACTGTGCAATCA---CTGAAATTG
 DR25006 GAAACCTGAGGACACGGCCATGTATTACTGCGCGGTAGATGGTTGGACCC
 DR20006 CAAACCTGAGGACACGGCCATGTATTACTGTGCG-----CC
 DR21006 GAACCCGGAGGACACTGCCGACTACTACTGCGCTGCAAATCAATTAGC--
 DR09006 GAGGCCTGAGGACACGGCCGTGTATTACTGTGCGGCAGATTG-----
 DR17006 GAAACCTGAGGACACGGCCATCTATTACTGTGCGGCAG-----CGGACCC
 DR13006 GCAACCTGACGACACTGGCGTGTACTACTGTGCGGCC-----CAA
 DR02006 GAAACCCGAGGACACGGCCGTGTATTACTGTGGGGCAGT-----

DR01006 -----A-AC--CTGGGGGTTATTGTGGGTA-
 DR27006 -----GTAC--CCGTGCCATCTCCTTGATG-
 DR03006 ACTTGGGACCT-----GGCG-CCATT-----CTTGATTG
 DR11006 AGATGGGGGCTTGTGATGCGAGATGGGCGACCTTAGC--GACAAGGAC-G
 DR24006 AGATGGGGGCTTGTGATGCGAGATGGGCGACCTTAGC--GACAAGGAC-G
 DR16006 GCGTTTTT-CTAGTCCTGTGGGAGCACTTC-TAGAC---TCGAAAGTAG
 DR19006 TAGTTTTTACTGGTACT-----GCAC-----C---ACG-----G
 DR07006 GGTGGTAGTTGTGAATTGC---CTTTGC-----TATTTGACTA-----
 DR16006 TGTTACATGCCGACTATGC---CCGCTCCCCGATACGAGACAGTTTGG
 DR20006 AGTGGTATGGGTGCAATTT---AAGGACTACTTTTACT---C-----G
 DR25006 GGAAGGAAG--GGGGAATCGGGTTAC---CTGGTTCGGTCCAATGTGAA
 DR20006 GGTGAA-----TATC-----CTATTGCAGAC--ATGTGTT
 DR21006 ---TGGTGGCTGGTATT-----TGGACCCGAATTACTGG-CTCTCTGTG
 DR09006 ---GAAATACTGGA---CTTGTGGTGC--CCAGA-CTGG-----AG
 DR17006 AAGTATATATTATAGTATC-----CTCCNNAT-----
 DR13006 AAGAAGGATCGTA-----CTAGATGGGC-----CGAGCCT-----
 DR02006 -----CTCCCTAA--TGGACCGAATTTC

DR01006 --TGGGTANTGCCTCTGGGGCCAGGGGACCCAGGTCACCGTCTCCTCACT
 DR27006 --T-----CTGGGGCCAGGGGACCCAGGTCACCGTCTCCTCACT
 DR03006 AAAAAGTATAAGTACTGGGGCCAGGGGACCCAGGTCACCGTCTCCTCACT
 DR11006 TTTGCGTATAACTACTGGGGCCGGGGGACCCAGGTCACCGTCTCCTCACT
 DR24006 TTTGCGTATAACTACTGGGGCCGGGGGACCCAGGTCACCGTCTCCTCACT
 DR16006 CGA-CT-ATAACTATTGGGGCCAGGGGATCCAGGTCACCGTCTCCTCACT
 DR19006 CGC-CTTATAACGTCTGGGGTCAGGGGACCCAGGTCACCGTCTCCTCACT
 DR07006 CTGGG-----GCCAGGGCACCAGGTCACCGTCTCCTCACT
 DR16006 CTGGGATGATTTT-----GGCCAGGGGACCCAGGTCACCGTCTCCTCACT
 DR20006 CTGGG-----GCCAGGGGACCCAGGTCACCGTCTCCTCACT
 DR25006 GATGGTTATAACTATTGGGGCCAGGGGACCCAGGTCACCGTCTCCTCAC-
 DR20006 CGAGAT----ACG---GCGACCCGGGGACCCAGGTCACCGTCTCCTCAC-
 DR21006 GGTGCATATGCCATCTGGGGCCAGGGGACCCAGGTCACCGTCTCCTCAC-
 DR09006 GATACTTCGGACAG-TGGGGTCAGGGGGCCAGGTCACCGTCTCCTCACT
 DR17006 --TGAGTATAAGTACTGGGGCCAGGGGACCCAGGTCACCGTCTCCTCA--

FIGURE 7(3)
 SUBSTITUTE SHEET

```
DR13006 CGAGAATGGAACAAC TGGGGCCAGGGGACCCAGGTCACCGTCTCCTCA--
DR02006 CCAACATGGG--TGCCGGGGCCAGGGAACCCAGGTCACCGTCTCCT----

DR01006 AG----TTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR27006 AG----TTACCCGTACGAGCTTCCGGACTACGGTTCCTTAATAGAATTC
DR03006 AGCTAGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR11006 AG----TTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR24006 AGCTAGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR16006 ----AGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR19006 ----AGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR07006 ----AGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR16006 ----AGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR20006 ----AGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR25006 ---TAGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR20006 ---TAGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR21006 ---TAGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR090J6 AGCTAGTTACCCGTACGACGTTCCGGACTACGGTTCCTTAATAGAATTC
DR17006 -----
DR13006 -----
DR02006 -----TA
```

FIGURE 7(4)




FIGURE 8

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINE(S) OR MARK(S) ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.